

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/003061

International filing date: 22 March 2005 (22.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 04019293.2
Filing date: 13 August 2004 (13.08.2004)

Date of receipt at the International Bureau: 23 May 2005 (23.05.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

03. 05. 2005

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04019293.2

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 04019293.2
Demande no:

Anmeldetag:
Date of filing: 13.08.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Complex Biosystems GmbH
Im Neuenheimer Feld 584
69120 Heidelberg
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Prodrug linker

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K47/48

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PL PT RO SE SI SK TR LI

Prodrug Linker

EPO - Munich
60

Field

13. Aug. 2004

- 5 The present invention is directed to polymeric prodrugs having reversible linkages to amino groups of biologically active entities such as peptides, proteins, natural products or synthetic chemical compounds.

Background

10

Polymers used in drug delivery systems are of interest because the therapeutic efficacy of both many low molecular weight and macromolecular medicinal agents may be significantly improved by combining them with polymers.

- 15 For example, covalent modification of biological molecules with poly(ethylene glycol) (PEG) has been extensively studied since the late 1970s. So-called PEGylated proteins have shown improved therapeutic efficacy by increasing solubility, reducing immunogenicity, and increasing circulation half-life in vivo due to reduced renal clearance and proteolysis by enzymes (see, for example, Caliceti P., Veronese F.M.,
20 Adv. Drug Deliv. Rev. 2003, 55, 1261-1277).

- Many small molecule medicinal agents, like alkaloids and anti-tumor agents, show low solubility in aqueous fluids. One way to solubilize these small molecule compounds is to conjugate them to hydrophilic polymers. A variety of water-soluble
25 polymers, such as human serum albumin, dextran, lectins, poly(ethylene glycol), poly(styrene-co-maleic anhydride), poly(N-hydroxypropylmethacrylamide), poly(divinyl ether-co-maleic anhydride), hyaluronic acid have been described for this purpose.

- 30 Depot formulations generating long-acting release profiles have been described using non-covalent drug encapsulation. Typically, a drug is mixed with polymer material and processed in such fashion, that the drug is captured in the polymer interior.

Resulting polymer-protein aggregates may be shaped as microparticles or gels. Drug release occurs if the polymer swells or degradation of the polymer allows for diffusion of the drug to the exterior. Degradation processes may be autohydrolytic or enzyme-catalyzed. Marketed drugs based on non-covalent polymer encapsulation are Lupron Depot and Nutropin Depot. A disadvantage of this approach is that in order to prevent uncontrolled, burst-type release of the non-covalently bound drug, encapsulation has to be highly efficient, requiring strong van der Waals contacts, frequently mediated through hydrophobic moieties, and many conformationally sensitive therapeutics such as proteins or peptides are rendered dysfunctional during the encapsulation process and/or during subsequent storage. Furthermore, dependence of the release mechanism upon biodegradation may cause interpatient variability.

A major challenge in cancer therapy is to selectively target cytotoxic agents to tumor cells. A promising method to accumulate small molecule anticancer agents in tumor tissue and decrease undesirable side effects of these agents is the attachment of the cytotoxin to a macromolecular carrier. The passive targeting of polymeric drug conjugates to tumors is based on the so-called enhanced permeability and retention effect (EPR) as described by Matsumura, Y. and Maeda, H., in *Cancer Res.*, 1986, vol 6, pp 6387-6392. As a result, several polymer-drug conjugates have entered clinical trial as anticancer agents.

However, many medicinal agents are inactive or show decreased biological activity when a polymer is covalently conjugated to the drug molecule. In these cases, a prodrug approach may be applied in such a fashion that the medicinal agent is released from the polymeric carrier in vivo to regain its biological activity. Prodrugs may be defined as therapeutic agents that are inactive per se but are predictably transformed into active metabolites (see B. Testa, J.M: Mayer in *Hydrolysis in Drug and Prodrug Metabolism*, Wiley-VCH, 2003, page 4). The reduced biological activity of the prodrug as compared to the released drug is of advantage if a slow or controlled release of the drug is desired. In this case, a relatively large amount of prodrug may be administered without concomitant side effects and the risk of overdosing. Release of the drug occurs over time, thereby reducing the necessity of repeated and frequent administration of the drug.

Definitions established by IUPAC

(as given under <http://www.chem.qmul.ac.uk/iupac/medchem/> (accessed on 8 March 2004))

5 Prodrug

A prodrug is any compound that undergoes biotransformation before exhibiting its pharmacological effects. Prodrugs can thus be viewed as drugs containing specialized non-toxic protective groups used in a transient manner to alter or to eliminate undesirable properties in the parent molecule.

10

Double prodrug (or pro-prodrug)

A double prodrug is a biologically inactive molecule which is transformed *in vivo* in two steps (enzymatically and/or chemically) to the active species.

15

Carrier-linked prodrug (Carrier prodrug)

A carrier-linked prodrug is a prodrug that contains a temporary linkage of a given active substance with a transient carrier group that produces improved physicochemical or pharmacokinetic properties and that can be easily removed *in vivo*, usually by a hydrolytic cleavage. This is shown graphically in Fig. 1.

20

Cascade prodrug

A cascade prodrug is a prodrug for which the cleavage of the carrier group becomes effective only after unmasking an activating group. This is shown graphically in Fig.

25 2.

Bioprecursor prodrug

A bioprecursor prodrug is a prodrug that does not imply the linkage to a carrier group, but results from a molecular modification of the active principle itself. This modification generates a new compound, able to be transformed metabolically or chemically, the resulting compound being the active principle.

30

Biotransformation

Biotransformation is the chemical conversion of substances by living organisms or enzyme preparations.

- 5 This invention applies in particular to carrier prodrugs and cascade prodrugs. In polymeric prodrugs, the biologically active moieties are often linked to a polymeric carrier moiety by a labile bridge formed between the carrier moiety and a hydroxy, amino or carboxy group of the drug molecule.
- 10 Cleavage of a carrier prodrug generates a molecular entity (drug) of increased bioactivity and at least one side product, the carrier. This side product may be biologically inert (for instance PEG) or may have targeting properties (for instance antibodies). After cleavage, the bioactive entity will reveal at least one previously conjugated and thereby masked or protected functional group, and the presence of this
- 15 group typically contributes to the bioactivity. Numerous macromolecular prodrugs are described in the literature where the macromolecular carrier is linked via a labile ester group to the medicinal agent (e.g. Y. Luo, MR Ziebell, GD Prestwich, "A Hyaluronic Acid – Taxol Antitumor Bioconjugate Targeted to Cancer Cells", *Biomacromolecules* 2000, 1, 208-218, J Cheng et al, *Synthesis of Linear, beta-Cyclodextrin Based*
- 20 *Polymers and Their Camptothecin Conjugates*, *Bioconjugate Chem.* 2003, 14, 1007-1017, R. Bhatt et al, *Synthesis and in Vivo Antitumor Activity of Poly(L-glutamic acid) Conjugates of 20(S)-Camptothecin*, *J. Med. Chem.* 2003, 46, 190-193 ; R.B. Greenwald, A. Pendri, C.D. Conover, H. Zhao, Y.H. Choe, A. Martinez, K. Shum, S. Guan, *J. Med. Chem.*, 1999, 42, 3657-3667; B. Testa, J.M: Mayer in *Hydrolysis in*
- 25 *Drug and Prodrug Metabolism*, Wiley-VCH, 2003, Chapter 8) In these cases, the conjugated functional group of the bioactive entity is a hydroxyl group or a carboxylic acid.
- Especially for biomacromolecules but also for small molecule polymer prodrugs, it
- 30 may be desirable to link the macromolecular carrier to amino groups (i.e. N-terminus or lysine amino groups of proteins) of the bioactive entity, as is shown in Fig. 3. This will be the case if masking the drug's bioactivity requires conjugation of a certain amino group of the bioactive entity, for instance an amino group located in an active center or a region or epitope involved in receptor binding. Also, during preparation of

the prodrug, amino groups may be more chemoselectively addressed and serve as a better handle for conjugating carrier and drug because of their greater nucleophilicity as compared to hydroxylic or phenolic groups. This is particularly true for proteins which may contain a great variety of different reactive functionalities. In this case
5 non-selective conjugation reactions lead to undesired product mixtures which require extensive characterization or purification and may decrease reaction yield and therapeutic efficiency of the product.

Prodrug activation may occur by enzymatic or non-enzymatic cleavage of the labile
10 bridge between the carrier and the drug molecule, or a sequential combination of both, i.e. an enzymatic step followed by a nonenzymatic rearrangement.

In an enzyme-free in vitro environment such as an aqueous buffer solution, almost any prodrug will undergo hydrolysis if labile linkages such as ester or amide bonds
15 are present in the molecule. This rate of hydrolysis may be very slow and not therapeutically useful. In an in vivo environment, esterases or amidases are typically present and may cause significant catalytic acceleration of the kinetics of hydrolysis from twofold up to several orders of magnitude (see, for example, R.B. Greenwald et al. J.Med.Chem. 1999, 42 (18), 3857-3867).

20 The labile bond may be engineered for enzyme selectivity for organ or cell-type targeting. Targeted release of the bioactive entity is effected, if an enzyme, that selectively cleaves the linkage, is specifically present in the organ or cell type chosen for treatment. Several examples have been published for the prodrug activation of
25 amine containing biologically active moiety by specific enzymes. In these cases, targeted release was the objective. G. Cavallaro et al., Bioconjugate Chem. 2001, 12, 143-151 described the enzymatic release of an antitumoral agent by the protease plasmin. Cytarabin was coupled via the tripeptide sequence D-Val-Leu-Lys to the polymer alpha, beta-poly(N-hydroxyethyl)-DL-aspartamide (PHEA). Enzymatic
30 release of cytarabin was effected by the protease plasmin which concentration is relatively high in various kinds of tumor mass.

Further examples for antitumoral polymeric prodrugs activated by specific enzymes like beta lactamase (R. Satchi-Fainaro et al., Bioconjugate Chem. 2003, 14, 797-804)

- and cysteine proteases like cathepsin B (R. Duncan et al. J. Contr. Release 2001, 74, 135-146) have been described. Wiwattanapatapee et al. (2003) described a dendrimer prodrug for colonic delivery of 5-aminosalicylic acid. The drug molecule was conjugated by an azo bond to generation 3 PAMAM dendrimer. 5-aminosalicylic acid is released in the colon by a bacterial enzyme called azo reductase (W. R. Wiwattanapatapee, L. Lomlim, K. Saramunee, J. Controlled Release, 2003, 88: 1-9).
- A.J. Garman et al. (A.J. Garman, S.B. Kalindjan, FEBS Lett. 1987, 223 (2), 361-365 1987) used PEG5000-maleic anhydride for the reversible modification of amino groups in tissue-type plasminogen activator and urokinase. Regeneration of functional enzyme from PEG-uPA conjugate upon incubation at pH 7.4 buffer by cleavage of the maleamic acid linkage followed first order kinetics with a half-life of 6.1 h. The prodrug cleavage was not investigated in the presence of enzymes, and it can be expected – as explained above - that in vivo proteases will significantly contribute to the cleavage of the labile amide linkage. A further disadvantage of this linkage is the lack of stability of the conjugate at lower pH values that limits this approach to active agents stable at basic pH values as purification of the active agent polymer conjugate has to be performed under basic conditions.
- R.B. Greenwald, A. Pendri, C.D. Conover, H. Zhao, Y.H. Choe, A. Martinez, K. Shum, S. Guan, J. Med. Chem., 1999, 42, 3657-3667 & PCT Patent Application WO-A-99/30727 described a methodology for synthesizing poly(ethylene glycol) prodrugs of amino-containing small molecule compounds based on 1,4- or 1,6-benzyl elimination. In this approach the amino group of the drug molecule is linked via a carbamate group to a PEGylated benzyl moiety. The poly(ethylene glycol) was attached to the benzyl group by ester, carbonate, carbamate, or amide bonds. The release of PEG from the drug molecule relied on the enzymatic cleavage of these masking groups. The cleavage of the release-triggering masking group is followed in this approach by the classical and rapid 1,4- or 1,6-benzyl elimination. As a consequence the highly reactive and potentially toxic quinone methide or quinonimine methide is liberated.

This is an example for a cascade carrier prodrug where the masking group is identical with the carrier and is shown diagrammatically in Fig. 4.

This linker system was also used for releasable poly(ethylene glycol) conjugates of proteins (S. Lee, R.B. Greenwald et al. *Bioconj. Chem.* 2001, 12 (2), 163-169).

Lysozyme was used as model protein because it loses its activity when PEGylation takes place on the epsilon-amino group of lysine residues. Various amounts of PEG linker were conjugated to the protein. Regeneration of native protein from the PEG conjugates occurred by enzymatic cleavage in rat plasma or in non-physiological high pH buffer.

Greenwald et al. published in 2000 a poly(ethylene glycol) drug delivery system of amino-containing prodrugs based on trimethyl lock lactonization (R.B. Greenwald et al. *J. Med. Chem.* 2000, 43(3), 457-487; PCT Patent Application No. WO-A-02/089789). In this prodrug system substituted o-hydroxyphenyl-dimethylpropionic acid is coupled to amino groups of drug molecules by an amide bond. The hydroxy group is linked to PEG by an ester, carbonate, or carbamate group. The rate determining step in drug release is the enzymatic cleavage of these functional groups followed by fast amide cleavage by lactonization, liberating a potentially toxic aromatic lactone side product.

A common disadvantage of the linker systems described by Greenwald is the release of a potentially toxic aromatic small molecule side product after cleavage of these prodrug linker molecules.

More recently, R.B. Greenwald et al. (Greenwald et al. *J. Med. Chem.* 2004, 47, 726-734) described a PEG prodrug system based on bis-(N-2-hydroxyethyl)glycin amide (bicin amide) linker. In this system two PEG molecules are linked to a bicin molecule coupled to an amino group of the drug molecule. The first two steps in prodrug activation is the enzymatic cleavage of both PEG molecules. Different linkages between PEG and bicin are described resulting in different prodrug activation kinetics. The main disadvantage of this system is the slow hydrolysis rate of bicin amide conjugated to the drug molecule ($t_{1/2} = 3$ h in phosphate buffer) resulting in the release of a bicin-modified prodrug intermediate that may show different pharmacokinetic and pharmacodynamic properties compared to the parent drug molecule.

The in vivo release of stable linker-drug intermediates or potentially toxic side products can be avoided, if the carrier is not serving as a masking group and the activating group is coupled to the carrier by means of a stable bond. Antczak et al.
5 (Bioorg Med Chem 9 (2001) 2843-48) describe a reagent which forms the basis for a cascade prodrug system for amine-containing drug molecules. In this approach an antibody serves as carrier, a stable bond connects the antibody to an activating moiety, carrying an enzymatically cleavable masking group. Upon enzymatic removal of the masking group, a second labile bond cleaves and releases the drug compound, as
10 shown in Fig. 5.


A drawback of predominantly enzymatic cleavage is interpatient variability. Enzyme levels may differ significantly between individuals resulting in biological variation of prodrug activation by enzymatic cleavage. Enzyme levels may also vary depending on
15 the site of administration, for instance it is known that in the case of subcutaneous injection, certain areas of the body yield more predictable therapeutic effects than others. To reduce this unpredictable effect, non-enzymatic cleavage or intramolecular catalysis is of particular interest (see, for example, B. Testa, J.M: Mayer in Hydrolysis in Drug and Prodrug Metabolism, Wiley-VCH, 2003, page 5).

20 Using enzyme specificity to control both targeting and release kinetics is a compromise. Enzyme-dependent release results in interpatient variability. Targeting through enzyme selectivity is of lower efficiency than for instance targeting through antibodies and may be achieved by more reliable means for instance by introducing
25 certain structural features into the polymer moiety of the prodrug.


For these reasons, there is a need to provide novel linker and/or carrier technologies for forming polymeric prodrugs of amine containing active agents in order to overcome the limitations of the described polymeric prodrugs.
30

Detailed description of the invention

The present invention addresses the disadvantages described above. The invention provides for a polymeric prodrug of Formula Ia or Ib.




Formula Ia



Formula Ib

wherein Y_1 to Y_5 , R_1 to R_4 , T , X , W , Nu and Ar are defined below:

Native drug release is effected by a two step mechanism. The first step is the rate-determining cleavage of the masking group:




from the polymeric prodrug in vivo.

As described above, cleavage of the masking group may be mediated by an enzymatic or a non-enzymatic step, such as pH-dependent hydrolysis or intramolecular cyclization. In the preferred embodiment of the invention, cleavage is effected non-enzymatically by intramolecular cyclization or catalysis. The half-life of the cleavage kinetics in an aqueous buffer of pH 7.4 at 37 °C of the masking group

according to the present invention is preferably between 1 hour and 6 months, more preferably between 1 day and 3 months, and most preferably between 1 day and 2 months.


- 5 The second and final step in release of the regenerated native drug is the fast, spontaneous and irreversible so-called 1,4- or 1,6 or 1,(4+2p) (in which $p = 2, 3, 4$ or higher) elimination of the



moiety of the remaining polymeric prodrug of formula Ia or formula Ib, respectively.

15

This mechanism of native drug release from a polymeric prodrug triggered by hydrolytic cleavage of the masking group followed by a 1,6-elimination step of the activating group is exemplified by an example of a polymeric prodrug according to the present invention.



Definition of Y₁ to Y₅, R₁ to R₄, T, X, W, Nu and Ar in formula Ia or Ib

T is D or A

and D is a residue of an amine-containing biologically active material including but
25 not limited to small molecule bioactive agents or biopolymers like proteins,
polypeptides and oligonucleotides (RNA, DNA), peptide nucleic acids (PNA),
and A is a leaving group.

Non-limiting examples of suitable leaving groups A include but are not limited to
30 chloride, bromide, fluoride, nitrophenoxy, imidazolyl, N-hydroxysuccinimidyl, N-
hydroxybenzotriazolyl, N-hydroxyazobenzotriazolyl, pentafluorophenoxy, N-
hydroxysulfosuccinimidyl, or any other leaving group known by those skilled in the
art.

Suitable organic small molecule bioactive moieties include, without limitation, moieties such as central nervous system-active agents, anti-infective, anti-neoplastic, antibacterial, anti-fungal, analgesic, contraceptive, anti-inflammatory, steroidal, vasodilating, vasoconstricting, and cardiovascular agents with at least one primary or secondary amino group. Non-exclusive examples of such compounds are

5 daunorubicin, doxorubicin, idarubicin, mitoxantron, aminoglutethimide, amantadine, diaphenylsulfon, ethambutol, sulfadiazin, sulfamerazin, sulfamethoxazol, sulfalen, clinafloxacin, moxifloxacin, ciprofloxacin, enoxacin, norfloxacin, neomycin B, sprectinomycin, kanamycin A, meropenem, dopamin, dobutamin, lisinopril, serotonin,

10 carbutamid, acivicin, etc.

Suitable proteins and polypeptides having at least one free amino group include but are not limited to ACTH, adenosine deaminase, agalsidase, albumin, alfa-1 antitrypsin (AAT), alfa-1 proteinase inhibitor (API), alteplase, anistreplase, ancrod serine

15 protease, antibodies (monoclonal or polyclonal, and fragments or fusions), antithrombin III, antitrypsins, aprotinin, asparaginases, biphalin, bone-morphogenic proteins, calcitonin (salmon), collagenase, DNase, endorphins, enfuvirtide, enkephalins, erythropoietins, factor VIIa, factor VIII, factor VIIa, factor IX, fibrinolysin, fusion proteins, follicle-stimulating hormones, granulocyte colony

20 stimulating factor (G-CSF), galactosidase, glucagon, glucocerebrosidase, granulocyte macrophage colony stimulating factor (GM-CSF), phospholipase-activating protein (PLAP), gonadotropin chorionic (hCG), hemoglobins, hepatitis B vaccines, hirudin, hyaluronidases, iduronidase, immune globulins, influenza vaccines, interleukins (1 alfa, 1 beta, 2, 3, 4, 6, 10, 11, 12), IL-1 receptor antagonist (rhIL-1ra), insulins,

25 interferons (alfa 2a, alfa 2b, alfa 2c, beta 1a, beta 1b, gamma 1a, gamma 1b), keratinocyte growth factor (KGF), transforming growth factors, lactase, leuprolide, levothyroxine, luteinizing hormone, lyme vaccine, natriuretic peptide, pancrelipase, papain, parathyroid hormone, PDGF, pepsin, platelet activating factor acetylhydrolase (PAF-AH), prolactin, protein C, octreotide, secretin, sermorelin, superoxide dismutase

30 (SOD), somatotropins (growth hormone), somatostatin, streptokinase, sucrase, tetanus toxin fragment, tilactase, thrombins, thymosin, thyroid stimulating hormone, thyrotropin, tumor necrosis factor (TNF), TNF receptor-IgG Fc, tissue plasminogen activator (tPA), TSH, urate oxidase, urokinase, vaccines, plant proteins such as lectins and ricins.

Also included herein is any synthetic polypeptide or any portion of a polypeptide with in vivo bioactivity. Furthermore, proteins prepared by recombinant DNA methodologies including mutant versions of aforementioned proteins, antibody
5 fragments, single chain binding proteins, catalytic antibodies and fusion proteins are included.

Preferred proteins are antibodies, calcitonin, G-CSF, GM-CSF, erythropoietins, hemoglobins, interleukins, insulins, interferons, SOD, somatropin, TNF, TNF-
10 receptor-IgG Fc.

X is a spacer moiety such as R5-Y6

Y₁, Y₂ can each be either O, S, or NR6, independently of each other.
15 Y₃, Y₅ can each be either O or S, independently of each other.
Y₄ is O, NR6, or -C(R7)(R8)-
Y₆ is O, S, NR6, succinimide, maleimide, unsaturated carbon-carbon bonds or any heteratom containing a free electron pair, or is not present.
R2 and R3 are selected independently from hydrogen, substituted or non-substituted
20 linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryl, cyano, nitro, halogen, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, etc.

The term "heteroalkyl" in the context of the present invention denotes (linear, cyclical
25 or branched) alkyl chains where the alkyl chains contain one or more heteroatoms, selected independently from O, S, N, P, Si, etc. or groups, selected independently from amide, ester, phosphonate ester, phosphate ester, double or triple bonds, carbamate, urea, thiourea, thiocarbamate, oxime, etc.

30 Each R4 substitution on Ar may be the same or different and is selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy, heteroaryloxy, etc.

R4 is selected preferably from small substituents such as hydrogen, methyl, ethyl, ethoxy, methoxy, and other C1 to C6 linear, cyclical or branched alkyls and heteroalkyls.

5

n is zero or a positive integer.

R7 and R8 are selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or
10 non-substituted heteroaryls, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, cyano, halogen, etc.

R5 is selected from substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryls,
15 carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, etc.

R6 is selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryls, etc.

20

R1 is a polymer.

Non-limiting examples for suitable polymers are polyalkyloxy-based polymers like poly(propylene glycol) or poly(ethylene glycol), dextran, chitosan, hyaluronic acid and derivatives, alginate, xylan, mannan, carrageenan, agarose, cellulose, starch,
25 hydroxyethyl starch (HES) and other carbohydrate-based polymers, poly(vinyl alcohols), poly(oxazolines), poly(anhydrides), poly(ortho esters), poly(carbonates), poly(urethanes), poly(acrylic acids), poly(acrylamides) such as poly(hydroxypropylmethacrylamide) (HMPA), poly(acrylates), poly(methacrylates) like poly(hydroxyethylmethacrylate), poly(organophosphazenes), poly(siloxanes),
30 poly(vinylpyrrolidone), poly(cyanoacrylates), poly(esters) such as poly(lactic acid) or poly(glycolic acids), poly(iminocarbonates), poly(amino acids) such as poly(glutamic acid), collagen, gelatin, copolymers, grafted copolymers, cross-linked polymers, hydrogels, and block copolymers from the above listed polymers.

Hydrogels may be defined as three-dimensional, hydrophilic or amphiphilic polymeric networks imbibing large quantities of water. The networks are composed of homopolymers or copolymers, are insoluble due to the presence of covalent chemical or physical (ionic, hydrophobic interactions, entanglements) crosslinks. The crosslinks provide the network structure and physical integrity. Hydrogels exhibit a thermodynamic compatibility with water which allows them to swell in aqueous media. (see.: N.A. Peppas, P. Bures, W. Leobandung, H. Ichikawa, Hydrogels in pharmaceutical formulations, Eur. J. Pharm. Biopharm. 2000, 50, 27-46). The chains of the network are connected in such a fashion that pores exist and that a substantial fraction of these pores are of dimensions of between 1 and 1000 nm. By selecting certain polymerization conditions, the hydrogel may be obtained in the form of an amorphous gel or as beaded resin. Such soft beads may have a diameter of between 1 and 1000 micrometer.

Hydrogels may be synthesized from the polymers and copolymers listed above and physically cross-linked or chemically cross-linked by radical, anionic or cationic polymerization, by chemical reactions like condensation or addition reactions as described in W.E. Hennink and C.F. van Nostrum, Adv. Drug Del. Rev. 2002, 54, 13-36.

Further examples include branched and hyperbranched polymers. Examples for such polymers include dendrimers and other dense star polymers. (R. Esfand, D.A. Tomalia, Drug Discov Today, 2001, 6(8), 427-436; P.M. Heegaard, U. Boas, Chem. Soc. Rev. 2004 (33(1), 43-63; S.M. Grayson, J.M. Frechet, Chem. Rev. 2001, 101 (12), 3819-3868).

R1 can also be a biopolymer like a protein. Non-limiting examples of such polymers include albumin, antibodies, fibrin, casein, and other plasma proteins.

Each R1 polymer can carry one or more biologically active substances linked to the polymer by the prodrug linker described herein or any other linker known to the person skilled in the art. The polymers may have further substituents and may be functionalized for attachment to the spacer moiety X. Non-limiting examples of such functional groups comprise carboxylic acid and activated derivatives, amino,

maleimide, thiol, sulfonic acid and derivatives, carbonate and derivatives, carbamate and derivatives, hydroxyl, aldehyde, ketone, hydrazine, isocyanate, isothiocyanate, phosphoric acid and derivatives, phosphonic acid and derivatives, haloacetyl, alkyl halides, acryloyl, arylating agents like aryl fluorides, hydroxylamine, disulfides like
5 pyridyl disulfide, vinyl sulfone, vinyl ketone, diazoalkanes, diazoacetyl compounds, epoxide, oxirane, and aziridine.

Preferred functional groups for the R1 polymer include but are not limited to thiol, maleimide, amino, carboxylic acid and derivatives, carbonate and derivatives,
10 carbamate and derivatives, aldehyde, and haloacetyl.

Especially preferred functional groups include thiol, maleimide, amino, carboxylic acid and derivatives, carbamate and derivatives, and carbonate and derivatives thereof.

15 Non-limiting examples for suitable bonds or groups formed between X and R1 include disulfide, S-succinimido, amide, amino, carboxylic ester, sulfonamide, carbamate, carbonate, ether, oxime, hydrazone, urea, thiourea, phosphate, phosphonate, etc.


20 Preferred bonds or groups formed between X and R1 comprise S-succinimido, amide, carbamate, and urea.

Preferably, the R1 polymers are well hydrated, degradable or excretable, nontoxic and
25 non-immunogenic in mammals. Preferred R1 polymers include polyalkoxy-based polymers like polyethylene glycol and polyethylene glycol reagents as those described in Nektar Inc. 2003 catalog "Nektar Molecule Engineering – Polyethylene Glycol and Derivatives for Advanced PEGylation" and branched, hyperbranched, cross-linked polymers and hydrogels.

30 W is selected from substituted or non-substituted linear, branched or cyclical alkyl, aryls, substituted aryls, substituted or non-substituted linear, branched or cyclical heteroalkyl, substituted or nonsubstituted heteroaryls, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or

cyclical heteroalkyloxy, aryloxy, heteroaryloxycyano, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, etc.

- W is selected preferably from non-toxic substituted or non-substituted linear,
5 branched or cyclical alkyls or heteroalkyls.



At least one Nu is present in Nu-W.

- Nu is a nucleophile that can perform a nucleophilic attack at the carbonyl carbon of
15 $Y_4 \overset{Y_1}{\parallel} Y_2$ and thus catalyse the cleavage of the masking group by intramolecular
catalysis or cyclization. Nucleophiles are any functional groups or heteroatoms
carrying at least one free electron pair. Preferred nucleophiles include primary,
secondary and tertiary amino groups, thiol, carboxylic acid, hydroxylamine,
hydrazine. In order to effectively catalyse the cleavage of the masking group, the
20 spacing between the nucleophile Nu and Y_2 is preferably between three and fifteen
atoms. More preferably, the spacing between Nu and Y_2 is between four and eight
atoms. The at least one nucleophile Nu may be attached anywhere to W (e.g. at the
terminus or in the middle of W) or may be part of W.

- 25 An especially preferred variation of $Y_4 \overset{Y_1}{\parallel} Y_2$ is $NR6 \text{---} \text{C}(=O) \text{---} O$ and R6 may also
be Nu-W and is not hydrogen. In this case, preferred variations of each Nu-W are
selected independently from substructures according to the following substructure:



R9, R10, R11, and R12 are selected independently from hydrogen, substituted or non-
substituted alkyl or heteroalkyl, substituted or non-substituted aryl or heteroaryl.


Preferably, R9, R10, R11, and R12 are selected independently from hydrogen, substituted or non-substituted alkyl.

Figure 8 shows an example according to formula Ia or Ib wherein the cleavage of the
5 masking group is by intramolecular cyclization.


In cases where Nu only catalyses the cleavage of the masking group by intramolecular catalysis, no cyclical product of the masking group is formed.

Surprisingly it was found, that the masking group can modify irreversibly the amine
10 containing biologically active moiety when the nucleophile Nu is absent in the masking group. As shown in the example section, during release of the bioactive moiety insulin from a polymeric prodrug with a pentanoyl masking group which is not part of the present invention (as it does not contain a nucleotide), approximately 30 % of the insulin molecule was modified with the masking group by acyl transfer. The
15 mechanism of an example of this modification where D contains an additional free amino group that serve as nucleophile for acyl transfer is shown in figure 9.

Ar of formula Ia or Ib is a multi-substituted aromatic hydrocarbon or a multi-substituted aromatic heterocycle. To be aromatic, the number of pi electrons must
20 satisfy the Hückel rule ($4n+2$) and the cycle has to be planar. A huge variety of compounds satisfy these criteria and thus are suitable as Ar in formula Ia or Ib. Non-limiting preferred aromatic moieties include:



wherein W is O, N, or S, independent from each other.



formula Ib have to be arranged on the aromatic ring in such a fashion that a 1,4- or 1,6- or 1,(4+2p), with p= 2, 3, 4 and higher, elimination can take place (see above).

30 For example, in the case of a 6-membered ring, the substituents have to be arranged *ortho* or *para*.

Preferred moieties for Ar are mono- and dicyclic aromatic hydrocarbons or aromatic heterocycles.

Especially preferred moieties are monocyclic five- or six-membered aromatic hydrocarbons or aromatic heterocycles.

5 General synthesis procedures of the polymeric prodrugs

Synthesis of representative examples of polymeric prodrugs according to the present invention is described in the Examples section.

10 Prodrugs of the present invention can be prepared in various different fashions. Figure 10 shows general routes for the synthesis of the polymeric prodrugs of the present invention according to formula Ia.

In a first method, intermediate (III) is provided by acylating Y_2 of starting material

(II) with
$$\text{Nu}-\text{W}-\text{Y}_4-\text{C}(=\text{O})-\text{Y}_1$$

15

For this, X or Nu may have to be protected with a reversible protecting group PG_1 .

Suitable protecting groups are described in TW Greene, P.G.M. Wuts, Protective groups in organic synthesis, 1999, John Wiley & Sons, 3rd ed.. From intermediate (III)

20 two alternative routes can be used to yield (Iaa). In a first route intermediate (III) is activated by an activating agent such as 4-nitrophenyl chloroformate or disuccinyl carbonate to yield (IV). The amine containing drug molecule is attached to (IV) to yield (V) by displacing the leaving group of the activated intermediate (IV). After deprotection of X such as by treating intermediate (V) with reagents like trifluoroacetic acid or DTT (where applicable) deprotected intermediate (V) is then
25 reacted with polymer R1 to yield the polymeric prodrug (Iaa).

In a second route the polymer R1 is first attached to the intermediate (III) after deprotection of X (where applicable) to form intermediate (VI). After an activation step intermediate (VII) is formed. (VII) is reacted with the amine containing drug
30 molecule to form the polymeric prodrug (Iaa).


In a second method, intermediate (VIII) is provided by activating starting material (II) by an activating agent such as 4-nitrophenyl chloroformate. For this, Y_2 and/or X may have to be protected with a protecting group PG_2 and/or PG_1 . Amine containing drug

is reacted with intermediate (VIII) to form (IX). In a first route, Y₂ of (IX) is selectively deprotected and acylated to form intermediate (V) which is further processed to (Iaa) as described above. In a second route X is selectively deprotected and reacted with polymer R1 to form intermediate (X). Y₂ of (X) is then deprotected and acylated to form the polymeric prodrug (Iaa).

In a third method starting material (II) is reacted with polymer R1 to form intermediate (XI). In one route, intermediate (XI) can be acylated to form intermediate (VI) which processed as described above to form polymeric prodrug (Iaa). In a second route, Y₂ is protected by the protecting group PG₂, activated and reacted with the amine containing drug molecule to form (X). Intermediate (X) is then processed as described above to form the polymeric prodrug (Iaa).

For all methods described, further functional groups such as Y₃ or nucleophiles present in Nu-W may have to be protected with suitable protecting groups.

Polymeric prodrugs according to formula Ib can be prepared by methods described above for prodrugs according to formula Ia using starting material IIb instead of II in figure 10.



It is understood, that linker structures according to the outlined invention and carrying protecting groups or leaving groups as described and used in the synthesis of corresponding polymeric prodrugs are considered within the range of the invention.

Application of the polymeric prodrugs in molecular therapy

30 One advantage of the described prodrug systems is to enable depot formulations without the need for encapsulation. Until now, many biocompatible materials like hydrogels with large pore sizes could not be used for depot formulations due to their lack of encapsulation properties. From such well-hydrated and mechanically soft biocompatible materials, biologically active moiety would be released too fast for most

therapeutic applications. In combination with the prodrug linkers described in this invention, the carrier material may be optimized for its biocompatibility properties as the release is solely governed by the linker cleavage kinetics and does not require chemical or enzymatic degradation of the polymer carrier itself.

5

A second advantage of the polymeric prodrugs of the present invention is their predominantly non-enzymatic cleavage: the half-life of the prodrug in suitably buffered human blood plasma of pH 7.4 (with aqueous buffer concentration < 50 %) is at least 50 % of the half-life of the prodrug in enzyme-free buffer pH 7.4.

10

This feature allows for better predictability and control of release rates after administration to a living organism and reduces interpatient variability.

Release rates are governed by a substantially non-enzymatic chemical reaction which is in turn dependent on the molecular structure of the linker. Systematic or random modifications of the chemical structure, for instance by changing substituents in one or more positions, for instance a masking group in a cascade prodrug, allows for the generation of prodrug linkers with differing release rates. It is therefore possible to create a variety of prodrug linkers and select those fast or slow cleaving prodrug linkers according to the demands posed by a given medicinal or therapeutic application.

20

One embodiment of the invention relates to such cascade-prodrugs where the removal of a masking group from an activating moiety triggers a release sequence.

25

Another advantageous feature which is part of this invention is the attachment of the polymer carrier through a stable covalent bond to an activating moiety involved in a double or cascade prodrug release mechanism. Typically in polymer pro-prodrugs, a bond between the polymer carrier and the activating moiety is cleaved, this being the release-controlling step, followed by a more rapid molecular rearrangement of the activating moiety and cleavage of a second labile linkage which is located between the activating moiety and the drug. After completion of this sequence, the polymer carrier as well as the activating moiety are released in addition to the drug. Due to the rearrangement, the activating moiety is released in a highly reactive form and may

30

cause direct damage to surrounding biomolecules, or potentially toxic derivatives of the activating moiety may be formed in vivo. As part of this invention, the activating moiety remains attached to the polymer carrier after drug release and therefore cannot diffuse into the environment. Polymer carrier attachment greatly reduces the reactivity
5 of the activating moiety and the probability of unwanted side effects.

In contrast to the enzymatic dependency of masking group removal as described by Antczak, C., Bauvois, B., Monneret, C., Florent J-C, Bioorg Med Chem 9, 2001, 2843), a higher level of control over release rates can be achieved if the masking
10 group has enzyme-independent, self-eliminating properties. The masking groups according to the present invention contain at least one nucleophile Nu. This nucleophile, which can be for example a primary, secondary or tertiary amino group can assist the cleavage of the masking group from the activating moiety by intramolecular catalysis or cyclization. For instance the masking group may contain
15 an amine group attached to a four carbon atom chain, which in turn is linked through an ester group to the activating moiety. The free amine may react with this ester in an intramolecular fashion, resulting in ester cleavage, amidation and cyclization of the masking group, forming a six-membered ring. Structural features of this masking group such as nucleophilicity of the amine group and ring-forming capacity may be
20 systematically optimized in order to precisely adjust the rate of prodrug cleavage. Such intramolecular reactions resulting in unmasking and subsequent rearrangement are highly independent from enzymes due to the fact that intramolecular reactions are generally preferred over intermolecular reactions and are shown diagrammatically in Fig.6.

25 In another embodiment of the invention, independency of prodrug cleavage from enzyme levels is achieved by providing a prodrug containing a sterically demanding carrier group as is shown in Fig. 7.

30 Such encapsulation or sterical protection by the sterically demanding carrier group may be conferred by a branched, hyperbranched, crosslinked or self-assembled structure of the carrier polymer. Such polymers tend to form a densely packed molecular volume, as exemplified for instance in dendrimers, dense star polymers or bead-shaped nano- and microparticles or amorphous gels. If the linkage of the

polymer carrier to the drug is located in the interior of the polymer carrier, the linked drug will be efficiently encapsulated and protected from enzymatic attack. In this case, sterical hindrance by the polymer prevents enzymes from accessing and cleaving the labile linkage.

5

In yet another embodiment, enzyme-independent prodrug cleavage is achieved by combining an intramolecular self-eliminating masking group with an encapsulating hyperbranched or crosslinked or self-assembled carrier.

10 A further advantage of the present invention is the release of an unmodified biologically active moiety. In cases where the biologically active moiety contains further reactive functional groups like amino groups of lysine residues in proteins, an unwanted side reaction between the masking group and the biologically active moiety can occur. The reactive functional groups of the biologically active moiety may react
15 with the masking group, forming a stable covalent bond and resulting in the release of a modified biologically active moiety. This potential side reaction is shown schematically in figure 9. The occurrence of such side reactions is shown in the examples section using polymeric prodrugs which are not part of the present invention with simple masking groups like a pentanoyl residue without a nucleophile Nu present
20 in the masking group. This side reaction is suppressed using polymeric prodrugs according to the present invention with intramolecularly activated masking groups that contain nucleophiles Nu (see experimental section).

Description of the Figures

25

- Fig. 1 shows a carrier-linked prodrug.
Fig. 2 shows a cascade prodrug.
Fig. 3 shows cleavage of a prodrug with an amino group.
Fig. 4 shows prodrug cleavage from a carrier which is identical with a masking group.
30 Fig. 5 shows cleavage of a cascade prodrug system
Fig. 6 shows cleavage and intramolecular reaction of the masking group.
Fig. 7 shows cleavage of a prodrug having a sterically demanding carrier group.
Fig. 8 shows cleavage of the masker group by intramolecular cyclisation.
Fig. 9 shows a possible side reaction of polymeric prodrug activation

Fig.10 shows general synthesis methods

Fig.11 shows mass spectra of prodrug released insulin molecules

Examples

Materials

Fmoc-amino acids, resins and PyBOP were purchased from Novabiochem and are
5 named according to the catalogue. Fmoc-Ado-OH was obtained from Neosystem. All
additional chemicals were purchased from Sigma Aldrich. Recombinant human
insulin was from ICN Biomedicals (USA). Maleimide-PEG5k was obtained from
Nektar (USA). 5-(and-6)-carboxyfluorescein succinimidyl ester (mixed isomers) was
obtained from Molecular Probes.

Solid phase synthesis reaction medium

10 Solid phase synthesis was performed on NovaSyn TG Sieber amide resin with a
loading of 0.17 mmol/g or 2-chlorotrityl chloride resin with a loading of 1.4 mmol/g.
Syringes equipped with polypropylene frits were used as reaction vessels.

Standard coupling cycle for fmoc-protected amino acids

For fmoc protecting-group removal, the resin was repeatedly (three times, 4 min each)
agitated with 2/2/96 (v/v/v) piperidine/DBU/DMF and repeatedly (six times) washed
with DMF.

20 Coupling of fmoc-protected amino acids to free amino groups on resin was achieved
by agitating the resin with 3 equivalents (eq) of fmoc-amino acid, 3 eq PyBOP and 6
eq DIEA in relation to free amino groups in DMF for 60 min.
Finally, the resin was repeatedly (five times) washed with DMF.

Standard cleavage protocol for TentaGel Sieber amide resin

25 Upon completed synthesis, the resin was washed with DCM, dried in vacuo and
treated repeatedly (five times) with 97/2/1 (v/v) DCM/TES/TFA. After evaporation,
compounds were purified by preparative RP-HPLC (Waters 600).

Standard cleavage protocol for 2-chlorotrityl chloride resin


30 Upon completed synthesis, the resin was washed with DCM, dried in vacuo and
treated two times for 30 minutes with 65/35 (v/v) HFIP/DCM. After combining the
eluates the volatile components were evaporated.

Analysis

Mass spectrometry (MS) was performed on a Waters ZQ 4000 ESI instrument and spectra were, if necessary, interpreted by Waters software MaxEnt.

- 5 Size exclusion chromatography was performed using an Amersham Bioscience AEKTAbasic system equipped with a Superdex 200 column (Amersham Bioscience).
NMR spectra were recorded on a Bruker AC300.


Overview - synthesis of polymeric prodrugs according to formula Ia with ester-linked masking groups




Synthesis of polymeric prodrugs according to formula Ia with carbamate-linked masking groups

a)

5




b)




Synthesis of polymeric prodrugs according to formula Ib with a carbamate-linked masking group


5 a)



b)



Synthesis of compound 2




5

2

Mmt-chloride (1 eq) and mercaptopropionic acid (1.1 eq) were dissolved in TFA and incubated for 30 min. The solvent was removed under reduced pressure. The product was dissolved in pyridine, diluted in water, acidified by acetic acid and extracted with
10 ether. The ether phase was separated and dried over Na₂SO₄. Solvent was removed under reduced pressure and product **2** was purified by RP-HPLC.

Synthesis of compounds 3a and 3b

15



3a R1 = H

20 **3b** R1 = OMe

Octopamine hydrochloride **1a** or normetanephrine hydrochloride **1b** (2 eq), DIEA (4 eq), and PyBOP (1 eq) were dissolved in DMF, **2** (1eq) was added and the mixture was reacted for 50 min at room temperature. After addition of acetic acid (7 eq) products **3a** or **3b** were purified by RP-HPLC.


25

3a: MS [M+Na]⁺ = 536 (MW+Na calculated = 536.2 g/mol)

3b: MS [M+Na]⁺ = 566 (MW+Na calculated = 566.2 g/mol)

30

Synthesis of mercaptothiazolide 4




2-Mercaptothiazoline and triethylamine (1.5 eq) were dissolved in dry THF and pentanoyl chloride (1 eq) was added. The mixture was stirred for 1 h at 50 °C under
10 inert gas atmosphere and was allowed to cool to room temperature. 0.5 N aqueous HCl was added and the separated organic phases were dried over Na₂SO₄. After concentration *in vacuo* the residue was purified by silica gel column chromatography using heptane/ethyl acetate (1/1) as mobile phase. Mercaptothiazolide 4 was collected as a viscous yellow oil.

15

4 R_f (heptane/ethyl acetate 1:1) = 0.7

Synthesis of compounds 5a and 5b



25


General synthesis protocol:

1 g 2-chlorotrityl chloride resin (loading 1.6 mmol/g) was incubated for 1 h with 850 mg (2.4 mmol) Fmoc-Ile-OH and 840 µl (4.8 mmol) DIEA in 5 ml 1/1 DCM/DMF. After fmoc removal and washing of the resin with DMF, boc-aminobutyric acid or
30 boc-aminohexanoic acid, respectively, were coupled to 0.5 g resin each according to the standard coupling method. Compounds 5a or 5b, respectively, were cleaved from the resin with 97/1/2 (v/v) DCM/TFA/TES for 45 min. After neutralisation with pyridine, solvents were removed under reduced pressure and 5a or 5b were purified by RP-HPLC.

5a MS $[M+Na]^+ = 339.2$ (MW+Na calculated = 339.4 g/mol)

5b MS $[M+Na]^+ = 367.4$ (MW+Na calculated = 367.5 g/mol)

5 Synthesis of compound 6a




Mercaptothiazolide 4 (1 eq), phenol 3a (4 eq) and DMAP (4 eq) were refluxed in DCM for 2 h under nitrogen atmosphere. After neutralization with acetic acid, the solvent was removed *in vacuo* and product 6a was purified by RP-HPLC.

15

6a MS $[M+Na]^+ = 620$ (MW+Na calculated = 620.3 g/mol)

Synthesis of compounds 6b to 6e



General synthesis protocol:

6d	R1 = OMe
6e	R1 = H

20 Carboxylic acid 5a, 5b or Z-Lys(Boc)-OH (1 eq), phenol 3b (1 eq), DIC (1 eq), and DMAP (2 eq) in DMF were reacted for 1h at room temperature. After addition of acetic acid (4 eq) the resulting carboxylic esters 6c, 6d or 6b, respectively, were purified by RP-HPLC.

6e was synthesized as described above using 5b and 3a as starting materials.

25

6b MS $[M+Na]^+ = 928$ (MW+Na calculated = 928.6 g/mol)


6c MS $[M+Na]^+ = 864$ (MW+Na calculated = 864.5 g/mol)

6d MS $[M+Na]^+ = 892$ (MW+Na calculated = 892.6 g/mol)

6e MS $[M+Na]^+ = 862$ (MW+Na calculated = 862.6 g/mol)

30

Synthesis of compounds 7a to 7e




General synthesis protocol:

Alcohols **6a**, **6b**, **6c**, **6d**, or **6e** (1 eq), 4-nitrophenyl chloroformate (10 eq), and DIEA (10 eq) were stirred in dry dioxane for 3 h at room temperature under nitrogen atmosphere. After addition of acetic acid (25 eq) the mixtures were diluted with 7/3 (v/v) acetonitrile/H₂O and the carbonates **7a**, **7b**, **7c**, **7d**, or **7e**, respectively, were purified by RP-HPLC.

7a MS [M+Na]⁺ = 785 (MW+Na calculated = 785.5 g/mol)
7b MS [M+Na]⁺ = 1093 (MW+Na calculated = 1093.7 g/mol)
7c MS [M+Na]⁺ = 1029 (MW+Na calculated = 1029.6 g/mol)
7d MS [M+Na]⁺ = 1057 (MW+Na calculated = 1057.6 g/mol)
7e MS [M+Na]⁺ = 1027 (MW+Na calculated = 1027.6 g/mol)

Synthesis of compounds 8a to 8c (N^αAl-linker-insulin)



General synthesis protocol:


Rh-Insulin in 1/1 (v/v) DMSO/DMF was mixed with a solution of 0.9 eq carbonate **7a**, **7c**, or **7d** in DMSO. The resulting solutions were adjusted to basic pH with DIEA and stirred for 1.5 h at RT. RP-HPLC purification gave Mmt-protected intermediates. After lyophilization, the Mmt-protected intermediates were mixed with 95:5 (v/v) TFA/triethylsilane and stirred for 5 min. Volatiles were removed under nitrogen flow and **8a**, **8b** or **8c**, respectively, were purified by RP-HPLC and lyophilized. Regioselectivity of insulin modification was verified by DTT reduction and MS analysis.

25 **8a MS** $[M+2H]^{2+} = 3078.9$; $[M+3H]^{3+} = 2053.2$ $[M+4H]^{4+} = 1540.6$ (MW calculated = 6158 g/mol)

8b MS $[M+2H]^{2+} = 3152.9$; $[M+3H]^{3+} = 2100.6$ $[M+4H]^{4+} = 1575.8$ (MW calculated = 6302 g/mol)

30 **8c MS:** $[M+3H]^{3+} = 2110.7$; $[M+4H]^{4+} = 1583.7$; $[M+5H]^{5+} = 1266.6$ (MW calculated = 6330 g/mol)

Synthesis of compounds 8d to 8g (N^{εB29}-Fluorescein-N^{αA1}-linker-insulin)



Synthesis of N^{εB29}-fluorescein insulin:

80 mg (13.8 μmol) rh-insulin were dissolved in 4 ml 1/1 (v/v) DMF/DMSO and 40 μl
DIEA were added. 8 mg (17 μmol) 5-(and-6)-carboxyfluorescein succinimidyl ester
were added and the solution was stirred for 30 min at room temperature. 4 ml 5/5/1
(v/v/v) acetonitrile/water/acetic acid were added, product N^{εB29}-fluorescein insulin
was purified by RP-HPLC and lyophilized. The conjugation site was verified by
reduction of N^{εB29}-fluorescein insulin with 1,4-dithiothreitol, protease digestion and
MS analysis.

MS: [M+2H]²⁺ = 3084.0; [M+3H]³⁺ = 2054.6 (MW calculated = 6166 g/mol)

General synthesis protocol:

N^{B29}-fluorescein insulin in 1/1 (v/v) DMF/DMSO was mixed with a solution of 0.9 eq carbonate **7b**, **7c**, **7d**, or **7e** in DMSO. The resulting solutions were adjusted to basic pH with DIEA and stirred for 3 h at RT. RP-HPLC purification gave Mmt-protected intermediates.

After lyophilization, the intermediates were dissolved in 95/5 (v/v) TFA/triethylsilane and stirred for 5 min. Volatiles were removed under nitrogen flow and **8d**, **8e**, **8f**, or **8g** were purified by RP-HPLC and lyophilized.

- 10 **8d** MS: $[M+2H]^{2+} = 3364.1$; $[M+3H]^{3+} = 2242.7$; $[M+4H]^{4+} = 1681.5$ (MW calculated = 6724 g/mol)
8e MS: $[M+3H]^{3+} = 2219.2$ $[M+4H]^{4+} = 1665.9$; $[M+5H]^{5+} = 1332.8$ (MW calculated = 6660 g/mol)
8f MS: $[M+3H]^{3+} = 2229.7$ $[M+4H]^{4+} = 1673.3$; $[M+5H]^{5+} = 1337.7$ (MW calculated
15 = 6689 g/mol)
8g MS: $[M+3H]^{3+} = 2218.7$ $[M+4H]^{4+} = 1664.9$ (MW calculated = 6659 g/mol)


Synthesis of compounds 9a to 9g (mono-pegylated insulin compounds)

General synthesis protocol:

- 20 70 µl of a solution of **8a**, **8b**, **8c**, **8d**, **8e**, **8f**, or **8g** (500 µM) in 1/4 (v/v) acetonitrile/water were mixed with 7 µl of a solution of 10 mM maleimide-PEG5k in 1/4 (v/v) acetonitrile/water and 10 µl of 0.5 M sodium phosphate buffer (pH 7.0) and incubated for 15 min. Compounds **9a**, **9b**, **9c**, **9d**, **9e**, **9f**, or **9g**, respectively, were purified by SEC (column: Superdex 200, flow rate: 0.75 mL/min) using 10 mM
25 HEPES buffer (pH 7.4), 150 mM NaCl, 3 mM EDTA, and 0.005 % Tween as mobile phase. The collected eluates (approximately 1.5 mL) were directly used as such for release rate determination.

9a through 9g: SEC retention time: 19.5 min

Synthesis of compounds 11a and 11b



11a R1 = R2 = Me

11b R1 = Me, R2 = H

10 **General synthesis protocol:**

3,5-Dimethyl-4-hydroxy-acetophenone (5.0 mmol) **10a** or 4-hydroxy-3-methyl-acetophenone **10b** (0.75 g, 5.0 mmol) and CuBr₂ (1.7 g, 7.5 mmol) were dissolved in 10 ml ethyl acetate and refluxed for 2 h. Solid byproducts were removed by filtration. The filtrate was evaporated and crude products **11a** or **11b** were purified by RP-HPLC.


11a: Yield 754 mg (62 %)

MS [M+H]⁺ = 243.1/245.1 (MW+H calculated = 244.1 g/mol)

11b: Yield 533 mg (47 %)

20 MS [M+H]⁺ = 229.2/231.1 (MW+H calculated = 230.1 g/mol)

Synthesis of compounds 12a and 12b



12a R1 = R2 = Me

12b R1 = Me, R2 = H

30 **Synthesis protocol:**

500 mg **11a** (2.06 mmol) or 472 mg (2.06 mmol) **11b** and 576 mg (4.11 mmol) hexamethylenetetramine were dissolved in 20 ml of trichloromethane and refluxed for 30 min. The solvent was removed in vacuo. 4 ml ethanol and 2 ml of concentrated HCl were added and the slurry was heated to 50 °C for 4 h. The mixture was

concentrated in vacuo, diluted with acetonitrile/water and **12a** or **12b** were purified by RP-HPLC.

12a: Yield 547 mg (81 %) as TFA-salt


5 MS $[M+Na]^+ = 202.2$ (MW+Na calculated = 202.2 g/mol)

12b: Yield 455 mg (70 %) as TFA-salt

MS $[M+Na]^+ = 188.2$ (MW+Na calculated = 188,2 g/mol)

Synthesis of compound 13

10



13


15 500 mg (1.71 mmol) **12a** (TFA salt) were dissolved in 10 ml 1/1 (v/v) methanol/water, 129 mg (3.41 mmol) NaBH₄ were added and the mixture was stirred for 30 min at RT. 0.5 ml of acetic acid were added and **13** was purified by RP-HPLC.

13: Yield 313 mg (62 %) as TFA-salt

20 MS $[M+Na]^+ = 204.2$ (MW+Na calculated = 204.2 g/mol)

NMR (300 MHz, DMSO-d₆) δ [ppm] = 8.25 (s, 1H, Phenol), 7.84 (bs, 3H, NH₃⁺), 6.89 (s, 2H, CH_{ar}), 5.85 (d, 1H, Hydroxyl, J=3.7 Hz), 4.62 (m, 1H, CH_{Benzyl}), 2.93 (m, 1H, CH_a), 2.80 (m, 1H, CH_b), 2.17 (s, 6H, CH₃).

25 Synthesis of compounds 14a to 14c



30 **14a** R1 = R2 = Me, R7 = H

14b R1 = R2 = H, R7 = Me

14c R1 = OMe, R2 = H, R7 = Me

13 (TFA salt, 159 mg, 0.541 mmol), synephrine (335 mg, 2.00 mmol), or metanephrine (HCl salt, 281 mg, 1.20 mmol) were coupled to compound **2** as described for compound **3a** to yield **14a**, **14b**, or **14c**, respectively.

5 **14a**: Yield 254 mg (87 %)

MS $[M+Na]^+ = 564.7$ (MW+Na calculated = 564.3 g/mol)


14b: Yield 760 mg (72 %)

MS $[M+Na]^+ = 550.2$ (MW+Na calculated = 550.3 g/mol)

14c: Yield 530 mg (80 %)

10 MS $[M+Na]^+ = 580.4$ (MW+Na calculated = 580.4 g/mol)

Synthesis of compounds **15c**, **15d** and **15f**



General synthesis protocol:

1 g 2-chlorotriyl chloride resin (loading 1.4 mmol/g) was incubated for 1 h with
20 N,N'-dimethylpropane-1,3-diamine (for synthesis of **15c**), or N,N'-diethylpropane-
1,3-diamine (for synthesis of **15d**) or N,N'-dimethylethane-1,2-diamine (4 eq) (for
synthesis of **15f**) in DCM. After washing of the resin with DMF, amines were
acetylated with 1/1/2 (v/v/v) acetic anhydride/pyridine/DMF for 14 h. The resin was
washed with THF and dried. LiAlH₄ (1 M in THF, 4 eq) was added dropwise to the
25 suspended resin in THF. The resulting suspension was stirred for 3 h at 45°C under
nitrogen atmosphere. After cooling, aqueous Rochelle's salt solution was added and
the resin was separated and dried. Compounds were cleaved from resin with 2/1 (v/v)
HFIP/DCM (2 x 30 min). The volatile components were evaporated and the products
15c, **15d** or **15f** were used in the following steps without further purification.


30

15c MS $[M+H]^+ = 131.2$ (MW = 130.1 g/mol)

15d MS $[M+H]^+ = 159.2$ (MW = 158.1 g/mol)

15f MS $[M+H]^+ = 117.1$ (MW = 116 g/mol)

Synthesis of compounds **16a** to **16f**



5

16a R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl, R7 = H

16b R1 = OMe, R2 = H, R5 = Et, R6 = 2-(diethylamino)ethyl, R7 = H

16c R1 = OMe, R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

16d R1 = R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

10 **16e** R1 = OMe, R2 = H, R5 = Et, R6 = 3-(diethylamino)propyl, R7 = Me

16f R1 = R2 = H, R5 = Et, R6 = 3-(diethylamino)propyl, R7 = Me

14a (120 mg, 0.222 mmol) was dissolved in 1.5 ml of dry THF. p-Nitrophenyl-chloroformate (45 mg, 0.222 mmol) and DIEA (113 μ l, 0.665 mmol) were added and
15 the mixture was stirred for 30 min at RT. **15a** (N,N,N'-trimethyl-ethylene-1,2-diamine) (72 μ l, 0.554 mmol) was added and stirring was continued for 30 min. The solvent was removed in vacuo, 100 μ l of AcOH were added and **16a** was purified by RP-HPLC.

20 **16b** was synthesized as described above from **3b** (80 mg, 0.15 mmol) and **15b** (N,N,N'-triethyl-ethylene-1,2-diamine) (55 mg, 0.38 mmol).

16c or **16d** were synthesized as describe above from **14c** (56 mg, 0.1 mmol) or **14b** (53 mg, 0.1 mmol), respectively, and diamine **15c**.

16e or **16f** were synthesized as described above from **14c** (56 mg, 0.1 mmol) or **14b**,
25 respectively, (53 mg, 0.1 mmol) and diamine **15d**.

16a: Yield 120 mg (69 %) as TFA salt

MS $[M+Na]^+ = 692.4$ (MW+Na calculated = 692.9 g/mol)

16b: Yield 48 mg (40 %) as TFA salt

30 MS $[M+Na]^+ = 736.3$ (MW+Na calculated = 736.4 g/mol)

16c: Yield 8 mg (10 %) as TFA salt

MS $[M+Na]^+ = 736.4$ (MW+Na calculated = 736.4 g/mol)

16d: Yield 20 mg (25 %) as TFA salt

MS $[M+Na]^+ = 706.3$ (MW+Na calculated = 706.3 g/mol)

16e: Yield 2 mg (3 %) as TFA salt


MS $[M+Na]^+ = 764.6$ (MW+Na calculated = 764.4 g/mol)

16f: Yield 6 mg (8 %) as TFA salt

MS $[M+Na]^+ = 734.4$ (MW+Na calculated = 734.3 g/mol)

5

Synthesis of compound 17



10

17


Amine **12b** (TFA salt) was coupled to compound **2** as described for compound **3a**.

15 **17:** Yield 608 mg (74 %)

MS $[M+Na]^+ = 548.3$ (MW+Na calculated = 548.7 g/mol)

Synthesis of compounds 18a and 18b

20



18a R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl

18b R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl

25

383 mg (0.729 mmol) of **17** were reacted with p-nitrophenyl-chloroformate and N,N,N'-trimethyl-propane-1,3-diamine (**15e**) or **15f**, respectively, to yield **18a** or **18b** as described for compound **16a**.


30 **18a:** Yield 287 mg (50 %) as TFA salt

MS $[M+Na]^+ = 690.7$ (MW+Na calculated = 690,9 g/mol)

18b: Yield 148 mg (26 %) as TFA salt

MS $[M+Na]^+ = 690.9$ (MW+Na calculated = 690,9 g/mol)

Synthesis of compounds 16g and 16h



16g R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl

16h R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl

10 **18a** (287 mg, 0.367 mmol, TFA salt) was dissolved in 5 ml methanol, NaBH₄ (41 mg, 1.07 mmol) was added and the mixture was stirred for 30 min at RT. 0.5 ml of acetic acid were added and **16g** was purified by RP-HPLC.

18b (8 mg, 0.010 mmol, TFA salt) was reacted as described above to yield **16h**.

16g: Yield 201 mg (70 %) as TFA-salt


15 MS [M+Na]⁺ = 692.7 (MW+Na calculated = 692.9 g/mol)

16h: Yield 6 mg (77 %) as TFA-salt

MS [M+Na]⁺ = 692.7 (MW+Na calculated = 692.9 g/mol)

Synthesis of compounds 19a to 19h

20



25 **19a** R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl, R7 = H

19b R1 = OMe, R2 = H, R5 = Et, R6 = 2-(diethylamino)ethyl, R7 = H

19c R1 = OMe, R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

19d R1 = R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

19e R1 = OMe, R2 = H, R5 = Et, R6 = 3-(diethylamino)propyl, R7 = Me

30 **19f** R1 = R2 = H, R5 = Et, R6 = 3-(diethylamino)propyl, R7 = Me

19g R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl, R7 = H

19h R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl, R7 = H

Carbonates **19a** to **19h** were synthesized from **16a** to **16h**, respectively, as described for compound **7a**.

19a: Yield 98 mg (72 %) as TFA-salt

5 MS $[M+Na]^+ = 857.8$ (MW+Na calculated = 858.0 g/mol)

19b: Yield 6 mg (11 %) as TFA-salt

MS $[M+Na]^+ = 901.8$ (MW+Na calculated = 901.5 g/mol)

19c: Yield 1 mg (15 %) as TFA-salt

MS $[M+Na]^+ = 901.4$ (MW+Na calculated = 901.5 g/mol)

10 **19d:** Yield 8 mg (29 %) as TFA-salt

MS $[M+Na]^+ = 871.4$ (MW+Na calculated = 871.4 g/mol)

19e: Yield 0.3 mg (18 %) as TFA-salt

MS $[M+Na]^+ = 929.4$ (MW+Na calculated = 929.5 g/mol)

19f: Yield 4 mg (45 %) as TFA-salt

15 MS $[M+Na]^+ = 899.7$ (MW+Na calculated = 899.6 g/mol)

19g: Yield 6 mg (6 %) as TFA-salt


MS $[M+Na]^+ = 857.8$ (MW+Na calculated = 858.0 g/mol)

19h: Yield 0.8 mg (11 %) as TFA-salt

MS $[M+Na]^+ = 857.7$ (MW+Na calculated = 858.0 g/mol)

20

Synthesis of compounds **20a** to **20f**



30 **20a** R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl, R7 = H

20b R1 = OMe, R2 = H, R5 = Et, R6 = 2-(diethylamino)ethyl, R7 = H

20c R1 = OMe, R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

20d R1 = R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

20e R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl, R7 = H

20f R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl, R7 = H

Insulin derivatives **20a**, **20b**, **20c**, **20d**, **20e**, or **20f** were synthesized from **19a**, **19b**, **19c**, **19d**, **19g**, or **19h** respectively, as described for compound **8a**.

5

20a MS $[M+3H]^{3+} = 2077.3$ $[M+4H]^{4+} = 1559.2$ (MW calculated = 6231.3 g/mol)

20b MS $[M+3H]^{3+} = 2093.0$ $[M+4H]^{4+} = 1569.6$ (MW calculated = 6274 g/mol)


20c MS $[M+3H]^{3+} = 2090.8$ $[M+4H]^{4+} = 1568.7$ (MW calculated = 6274 g/mol)

20d MS $[M+3H]^{3+} = 2081.3$ $[M+4H]^{4+} = 1561.8$ (MW calculated = 6244 g/mol)

10 **20e** MS $[M+3H]^{3+} = 2077.1$ $[M+4H]^{4+} = 1558.2$ (MW calculated = 6231.3 g/mol)

20f MS $[M+3H]^{3+} = 2076.7$ $[M+4H]^{4+} = 1559.3$ (MW calculated = 6231.3 g/mol)

Synthesis of compounds **21a** to **21f** (mono-pegylated insulin derivatives)



21a R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl, R7 = H

21b R1 = OMe, R2 = H, R5 = Et, R6 = 2-(diethylamino)ethyl, R7 = H

21c R1 = OMe, R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

25 **21d** R1 = R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

21e R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl, R7 = H


21f R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl, R7 = H

Insulin derivatives **21a**, **21b**, **21c**, **21d**, **21e**, or **21f** were synthesized from compound

30 **20a**, **20b**, **20c**, **20d**, **20e**, or **20f**, respectively, as described for compound **9a**.

21a through **21f**: SEC retention time: 19.5 min

Synthesis of compounds 23a and 23b



23a R1 = Me, R2 = H

23b R1 = R2 = Me

o-Cresol (**22a**) or 2,6-dimethylphenol (**22b**) (1 eq), succinic anhydride (1 eq), and
10 AlCl₃ (3 eq) in nitrobenzene were heated to 100 °C for 1 h. The reaction mixture was
poured on HCl/ice and extracted with ether. The organic layer was extracted with 1 N
NaOH and the aqueous layer was acidified with concentrated HCl. The aqueous layer
was extracted with ether and the ether was evaporated. **23a** or **23b** were purified by
RP-HPLC.

15

23a: Yield 552 mg (31 %)


MS [M+Na]⁺ = 231.0 (MW+Na calculated = 231.2 g/mol)

NMR (300 MHz, DMSO-d₆) δ[ppm] = 12.05 (bs, 1H, CO₂H), 10.23 (s, 1 H, phenol
OH), 7.74 (s, 1H, CH_{ar}), 7.7 (d, 1H, CH_{ar}, ³J_{H,H} = 8.4 Hz), 6.86 (d, 1H, CH_{ar}, ³J_{H,H} =
20 8.4 Hz), 3.13 (t, 2H, C(O)CH₂, ³J_{H,H} = 6.4 Hz), 2.53 (t, 2H, CH₂CO₂, ³J_{H,H} = 6.4 Hz),
2.16 (s, 3H, CH₃)

23b: Yield 166 mg (15 %)

MS [M+Na]⁺ = 245.4 (MW+Na calculated = 245.2 g/mol)

25 Synthesis of compound 24




1.85 g (16.02 mmol) cysteamine hydrochloride were dissolved in 15 ml of TFA and
2.47 g (8.01 mmol) MmtCl were added. After stirring the mixture at RT for 20 min
30 the solvent was evaporated in vacuo. The residue was dissolved in diethyl ether and
extracted with saturated NaHCO₃, 1N H₂SO₄ and brine. The solvent was evaporated
and **24** was purified by RP-HPLC.

24: Yield 1.11 g (30 %) as TFA salt

TLC (AcOEt/Et₃N 99/1), R_f = 0.24

Synthesis of compounds 25a and 25b

5



25a R1 = Me, R2 = H

10 **25b** R1 = R2 = Me

23a or **23b** (1eq), HOBT (1.1 eq) and DIC (1 eq) were dissolved in DMF and stirred at RT for 30 min. **24** (TFA salt 1 eq) and DIEA (3 eq) were added and the solution was stirred for 60 min. Acetic acid was added and **25a** or **25b** was purified by RP-HPLC.

15

25a: Yield 552 mg (25 %)


MS [M+Na]⁺ = 562.7 (MW+Na calculated = 562.7 g/mol)

25b: Yield 15 mg (40 %)

20 MS [M+Na]⁺ = 576.6 (MW+Na calculated = 576.6 g/mol)

Synthesis of compounds 26a and 26b

25



26a R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

26b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl

267 mg (0.495 mmol) **25a** was reacted with p-nitrophenylchloroformate and N-(3-dimethylamino-propyl)-N',N'-dimethyl-propane-1,3-diamine (**15g**) to yield **26a** as described for compound **16a**.

30

26b was synthesized as described above using 15 mg **25b** and N,N,N'-trimethylethane-1,2-diamine (**15a**).

26a: Yield 282 mg (58 %) as double TFA salt


MS $[M+Na]^+ = 775.2$ (MW+Na calculated = 776.0 g/mol)

26b: Yield 17 mg (70 %) as TFA salt

MS $[M+Na]^+ = 704.5$ (MW+Na calculated = 704.6 g/mol)

5

Synthesis of compounds 27a and 27b



27a R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

27b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl

15 **26a** (272 mg, 0.277 mmol, double TFA salt) was dissolved in 5 ml methanol, $NaBH_4$ (42 mg, 1.09 mmol) was added and the mixture was stirred for 30 min at RT. 0.5 ml of acetic acid were added and **27a** was purified by RP-HPLC.

Alcohol **27b** was synthesized likewise from **26b** (17 mg, 25 μ mol, TFA salt).


20 **27a:** Yield 142 mg (52 %) as double TFA salt

MS $[M+Na]^+ = 777.9$ (MW+Na calculated = 778.0 g/mol)

27b: Yield 6 mg (40 %) as TFA salt

MS $[M+Na]^+ = 706.5$ (MW+Na calculated = 706.6 g/mol)

Synthesis of compounds 28a and 28b



28a R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

10 28b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl

Carbonates 28a or 28b were synthesized from 27a or 27b, respectively, as described for compound 7a.


15 28a: Yield 1 mg (29 %)

MS $[M+Na]^+ = 942.9$ (MW+Na calculated = 943.2 g/mol)

28b: Yield 1.5 mg (19 %)

MS $[M+Na]^+ = 871.6$ (MW+Na calculated = 871.7 g/mol)

Synthesis of compounds 29a and 29b



30 29a R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

29b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl

Insulin derivatives 29a or 29b were synthesized from 28a or 28b, respectively, as described for compound 8a.

29b MS $[M+3H]^{3+} = 2081.8$ $[M+4H]^{4+} = 1562.4$ (MW calculated = 6244 g/mol)

10




30b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl


20

30a and 30b: SEC retention time: 19.5 min


5




R2 =



R1 =



Synthesis of compound 31



31


5

31 was obtained according to the standard solid-phase synthesis protocol. The amino acids Fmoc-Dpr(Boc)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Ado-OH, and Fmoc-Dpr(Fmoc)-OH were coupled to NovaSyn TG Sieber amide resin.

- 10 After final fmoc removal the resin was agitated with 5 eq maleimidopropionic acid and 5 eq DIC in relation to amino groups in DMF for 30 min. **31** was cleaved from resin with TFA/TES/water 95/3/2 (v/v/v). After evaporation of solvent, product **31** was purified by RP-HPLC.

MS: $[M+H]^+ = 2494.6$ (MW calculated = 2495.4 g/mol)

Synthesis of compound 32



32


Compound **32** was obtained according to the standard solid-phase synthesis protocol.

- 5 The amino acids Fmoc-Cys(Mmt)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Ado-OH, and Fmoc-Dpr(Fmoc)-OH were coupled to NovaSynTG Sieber amide resin.

- After final fmoc removal the resin was agitated with 3 eq Boc-aminoxyacetic acid, 3 eq DIC, and 3 eq HOBt in relation to amino groups in DMF for 30 min. **32** was
- 10 cleaved from resin with DCM/TFA/TES 97/1/2 (v/v/v). After addition of 0.8 eq pyridine in relation to TFA the solvent was evaporated and product **32** was purified by RP-HPLC.

MS: $[M+H]^+ = 2688.2$ g/mol (MW calculated = 2688.8 g/mol)

Synthesis of compound 33



33

5

Compound 33 was obtained according to the standard solid-phase synthesis protocol. The amino acids Fmoc-Dpr(ivDde)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Lys(Fmoc)-OH, and Fmoc-Ado-OH were coupled to NovaSyn TG Sieber amide resin.


- 10 After final fmoc removal the resin was agitated with 3 eq 3,6,9-trioxadecanoic acid, 3 eq PyBOP, and 6 eq DIEA in relation to amino groups in DMF for 60 min.

To cleave the ivDde protecting group, the resin was treated three times with 2 % hydrazine in DMF. After washing, 3 eq Fmoc-Ser-OH was coupled with 3eq DIC and 3 eq HOBt for 30 min. After final fmoc removal resin was washed and the product

- 15 was cleaved from resin with DCM/TFA/TES 88/10/2 (v/v/v). Solvent was evaporated and the residue was oxidized with 10 eq sodium periodate in 3/2 (v/v) 0.1 M sodium phosphate pH 7/acetonitrile for 15 min to yield 33. Product 33 was purified by RP-HPLC and lyophilized.

MS: $[M+H]^+ = 3372.1 \text{ g/mol}$ (3372.8 g/mol)

Synthesis of compound 34



6 mg (2.4 μ mol) of compound **31** were dissolved in 1 ml 2/1 (v/v) acetonitrile/0.1 M sodium phosphate buffer pH 7 and 65 mg (24.2 μ mol) of compound **32** were added.

5 The solution was stirred at room temperature for 2 h and then the product purified by RP-HPLC and lyophilized (yield: 45 mg (78 %)).

The lyophilized product (45 mg) was dissolved in 0.5 ml DMF and 10 μ l DIEA were added. 5 mg (30 μ mol) 3-maleimidopropionic acid and 4.7 μ l (30 μ mol) DIC in 150 μ l
10 DMF were added and the reaction mixture was stirred at room temperature for 20 min, the product purified by RP-HPLC and lyophilized.

The lyophilized product was incubated for 10 min in 95/5 (v/v) TFA/water and then the solvent was removed in a stream of nitrogen. Product **34** was purified by RP-
15 HPLC and lyophilized (overall yield for all three steps: 20 mg (47 %)).

MS: 17700 – 18200 (broad peak) (MW calculated = 17749 g/mol)


Synthesis of compound **9h**

20 1.5 mg (225 nmol) **8g** and 5 mg (280 nmol) **34** were mixed, dissolved in 300 μ l 2/1 (v/v) 0.1 M sodium phosphate buffer pH 7/acetonitrile and incubated for 15 min at room temperature. The product was purified by RP-HPLC and lyophilized. (yield 4 mg, 160 nmol, 70 %)

25 The lyophilized product was dissolved in 200 μ l 0.1 M sodium citrate buffer pH 1.5 and 69 mg (20.5 μ mol) **33** in 200 μ l 2/1 (v/v) acetonitrile/sodium citrate buffer pH 1.5 were added. The mixture was stirred at room temperature for 24 h and product **9h** was purified by size exclusion chromatography (column: Superdex 200, buffer: 10 mM
30 HEPES pH 7.4, 0.005 % Tween-20, 3mM EDTA, flow rate: 0.75 ml/min)

SEC elution time: 15 min

Synthesis of compound 37a



250 mg (0.35 mmol) 2-chlorotrityl chloride resin (loading 1.4 mmol/g) was incubated
10 for 1.5 h with 308 mg (4 eq., 1.4 mmol) 4,7,10-trioxatridecane-1,13-diamine in 4 ml
DCM to yield 35a. The resin was washed with DCM and dried. 107 mg (0.7 mmol)
HOBt, 110 μ l (0.7 mmol) DIC, and 150 mg (0.9 mmol) 5-formyl salicylic acid in 3 ml
DMF were added and the resulting suspension was stirred for 1 h at RT to yield 36a.
After washing with DCM and THF, the resin was suspended in 6 mL THF and 3 ml (3
15 mmol) BH_3 THF (1 M in THF, 8.5 eq.) were added dropwise. The reaction mixture
was stirred for 18 h at 45°C under nitrogen atmosphere. After cooling 4 ml THF, 0.8
ml DIEA and 1.6 ml MeOH were added successively. 210 mg (0.84 mmol) I_2 (as a
concentrated THF solution) were added and the suspension was stirred for 1 h. The
resin was repeatedly washed (three times each) with THF, DMF, MeOH, and DCM.
20 The dried resin was reacted with 107 mg (0.7 mmol) HOBt, 110 μ l (0.7 mmol) DIC,
and 55 μ L (0.9 mmol) AcOH in 3 ml DMF for 1h. After washing of the resin with
DMF and DCM compound 37a was cleaved from resin with 2/1 (v/v) HFIP/DCM
(two times for 30 min). The volatile components were evaporated and the product 37a
was used in the following step without further purification.


25

37a: Yield 29 mg (20 %) as TFA salt

MS $[\text{M}+\text{Na}]^+ = 421.4$ (MW+Na calculated = 421.5 g/mol)

30

Synthesis of compound 38




24 mg (0.06 mmol) **37a**, 31 mg (0.06 mmol) PyBOP, 32 μ l (0.18 mmol) DIEA, and
10 23 mg (0.06 mmol) **2** in 0.5 mL DMF were reacted for 50 min at room temperature.
After addition of 50 μ l acetic acid product **38** was purified by RP-HPLC.

38: Yield 7 mg (15 %)

MS $[M+Na]^+ = 781.3$ (MW+Na calculated = 781.6 g/mol)

15

Synthesis of compound 39




20 300 mg (0.42 mmol) 2-chlorotrityl chloride resin (loading 1.4 mmol/g) was incubated
for 1.5 h with 245 mg (4 eq., 1.7 mmol) 1,8-diaminooctane in 4 ml DCM to yield **35b**.
The resin was washed with DCM and dried. 107 mg (0.7 mmol) HOBt, 110 μ l (0.7
mmol) DIC, and 150 mg (0.9 mmol) 5-formyl salicylic acid in 3 ml DMF were added
and the resulting suspension was stirred for 1 h at RT to yield **36b**. After washing with
25 DCM and THF, the resin was suspended in 6 mL THF and 3 ml (3 mmol) BH_3 THF
(1 M in THF) were added dropwise. The reaction mixture was stirred for 18 h at 45°C
under nitrogen atmosphere. After cooling 4 ml THF, 0.8 ml DIEA and 1.6 ml MeOH
were added successively. 210 mg (0.84 mmol) I_2 (as a concentrated THF solution)
were added and the suspension was stirred for 1 h. The resin was repeatedly washed

(three times each) with THF, DMF, MeOH, and DCM. The dried resin was reacted with 107 mg (0.7 mmol) HOBT, 110 μ l (0.7 mmol) DIC, and 55 μ L (0.9 mmol) AcOH in 3 ml DMF for 1h. After washing of the resin with DMF and DCM compound **37b**, 78 mg (0.39 mmol) p-nitrophenylchloroformate, and 210 μ l (1.2 mmol) DIEA in 1/1 (v/v) THF/DCM were reacted for 30 min at RT. The separated resin was suspended in 1/1 (v/v) THF/DCM and 210 μ l (1.2 mmol) N,N,N'-trimethylethylenediamine were added. The resulting suspension was stirred for 25 min at RT. The resin was separated and washed with DCM. Product **39** was cleaved from resin with 2/1 (v/v) HFIP/DCM (two times for 30 min). The volatile components were evaporated and the product **39** was purified by HPLC.

39: Yield 16 mg (8 %) as TFA salt

MS $[M+Na]^+ = 473.5$ (MW+Na calculated = 473.3 g/mol)

Synthesis of compound 40a




38 (7 mg, 9 μ mol) was dissolved in 200 μ l of dry THF. p-Nitrophenylchloroformate (2.0 mg, 10 μ mol) and DIEA (4.4 μ l, 25 μ mol) were added and the mixture was stirred for 30 min at RT. N,N,N'-Triethylethylenediamine (**15b**) (18 μ l, 0.1 mmol) was added and stirring was continued for 30 min. The solvent was removed *in vacuo*, 10 μ l of AcOH were added and **40a** was purified by RP-HPLC.

40a: Yield 1 mg (11 %) as TFA salt

MS $[M+Na]^+ = 951.1$ (MW+Na calculated = 951.8 g/mol)

Synthesis of compound 40b



5


15 mg (33 μ mol) **39**, 18 mg (33 μ mol) PyBOP, 23 μ l (0.13 mmol) DIEA, and 13 mg (35 μ mol) **2** in 0.5 mL DMF were reacted for 45 min at room temperature. After addition of 50 μ l acetic acid product **40** was purified by RP-HPLC.

10

40b: Yield 10 mg (37 %) as TFA salt

MS $[M+H]^+ = 811.5$ (MW+Na calculated = 810.5 g/mol)

Synthesis of compound 41a and 41b



Carbonate **41a** or **41b** was synthesized from **40a** or **40b** as described for compound **7a**


41a: Yield 0.4 mg as TFA salt

MS $[M+Na]^+ = 1116.8$ (MW+Na calculated = 1116.9 g/mol)

41b: Yield 2 mg (16 %) as TFA salt

MS $[M+H]^+ = 976.8$ (MW calculated = 975.8 g/mol)


Synthesis of compound 42



Insulin derivative **42** was synthesized from **41b** as described for compound **8a**.

42 MS $[M+3H]^{3+} = 2124.5$ $[M+4H]^{4+} = 1594.6$ (MW calculated = 6371 g/mol)


Synthesis of compound 43



Insulin derivative **43** was synthesized from **42** as described for compound **9a**.

43: SEC retention time: 18.0 min

Synthesis of rh-insulin loaded PEGA hydrogel 45



Maleimide derivatization of polyacrylamide based hydrogel (PEGA):

NH₂-PEGA hydrogel beads with 0.4 mmol/g loading and 150-300 μm bead size were purchased from Novabiochem.

2.5 g methanol-wet NH₂-PEGA-hydrogel (0.4 mmol/g NH₂-loading) was weighed into a syringe equipped with a polypropylene frit. Maleimide loading was adjusted by acylation employing a mixture of activated maleimidopropionic acid and acetic acid as described in the following. The hydrogel was washed 5 times with DMF and reacted with 13.5 mg (0.08 mmol) 3-maleimidopropionic acid, 115.2 μl (1.92 mmol)

acetic acid and 313 μ l (2 mmol) DIC in 4 ml DMF for 30 min. The maleimide derivatized hydrogel **44** was washed 10 times with DMF and DCM and finally with acetonitrile.


- 5 30 mg of maleimide derivatized resin **44** (loading 16 μ mol/g) was reacted with 3 mg of compound **20b** (480 nmol, 1.06 eq) in 600 μ l 20/80 (v/v) acetonitrile/50 mM phosphate buffer (pH 7.4) for 10 min to give rh-insulin loaded hydrogel **45**. The hydrogel **45** was washed 5 times with 50/50 (v/v) acetonitrile/water and three times with acetonitrile and dried under vacuum.

10

Synthesis of rh-insulin carbohydrate-based hydrogel **46**

15

20



46

- 25 NHS-activated "Sepharose 4 Fast Flow" hydrogel beads (chemically crosslinked agarose, crosslinker epichlorhydrin) were purchased from Amersham. 1.5 g ethanol-wet Sepharose hydrogel (150 mg dry hydrogel) was weighed into a syringe equipped with a polypropylene frit and reacted with 1 M 4,7,10-trioxatridecane-1,13-diamine in DMF for 30 min. After 5 washing steps with DMF, hydrogel was reacted with 8.5 mg (0.05 mmol) 3-maleimidopropionic acid, 57 μ l (0.95 mmol) acetic acid, 151 mg (1 mmol) HOBt and 158 μ l (1 mmol) DIC in 4 ml DMF for 30 min to give maleimide derivatized hydrogel. The hydrogel was washed 10 times with DMF and finally with acetonitrile.
- 30

1.5 mg **8c** was dissolved in 25/75 (v/v) acetonitrile/50 mM phosphate buffer pH 7.4 and reacted with 10.8 mg maleimide derivatized hydrogel for 10 min. The rh-insulin loaded hydrogel **46** was washed five times with 50/50 (v/v) acetonitrile/water and three times with acetonitrile and dried under vacuum.

5

Release of insulin or fluorescein-insulin from conjugates in buffer pH 7.4

Release of (fluorescein)-insulin from (fluorescein)-insulin conjugates **9a** to **9h**, **21a** to **21f**, **30a**, **30b**, and **43** was effected by linker hydrolysis in aqueous buffer pH 7.4.

10 Collected SEC eluates of (fluorescein) insulin conjugates (see above) were incubated at 37 °C and samples were taken at time intervals and analyzed by RP-HPLC (insulin conjugates) or SEC (fluorescein insulin conjugates) and UV detection. Peaks correlating with the retention time of native insulin or fluorescein-insulin, respectively, were integrated and plotted against incubation time, and curve-fitting
15 software was applied to estimate the corresponding halftime of release.

Release of insulin from hydrogel conjugates **45** and **46**

4 mg of **45** or 2 mg **46** were weighed into a test tube and incubated with 1 ml 10 mM
20 HEPES buffer pH 7.4, 150 mM NaCl, 0.005 % Tween at 37°C. 45 µl samples were taken at different time intervals and quantitatively analyzed for rh-insulin by a RP-HPLC assay. The rh-insulin peak was integrated and rh-insulin concentration was obtained from a standard curve. A first order release kinetic was fitted to the data points to give the linker half life.

25

MS-Analysis of released insulin from compound **9a**, **9b** and **30a**

Samples of buffer released insulin (see above) were analyzed by mass spectrometry. Figure 11 shows the mass spectra of released insulin from compound **9a**, **9b**, and **30a**.
30 The mass spectrum of insulin released from compound **9a** clearly shows a major side product (indicated by arrows), corresponding to the irreversibly pentanoyl-modified insulin. In this case, removal of the pentanoyl masking group was not by hydrolysis but by acyl transfer to the insulin. The mass spectrum of insulin released from compound **9b** and **30a** shows no modification.

Release of fluorescein-insulin from conjugate 9d and 9e in 80 % human plasma

Release of fluorescein-insulin from 9d or 9e was effected by hydrolysis in 80 % human plasma in 20 mM HEPES pH 7.4 at 37 °C. Samples were taken at time intervals and analyzed by SEC and UV detection. Peaks correlating with the retention time of fluorescein-insulin were integrated and plotted against incubation time, and curve-fitting software was applied to estimate the corresponding halftime of release.

Table: Polymeric prodrug hydrolysis

compound	t _{1/2} buffer pH 7.4	t _{1/2} human plasma
9a	40 h	nd
9b	55 h	nd
9c	4.5 d	nd
9d	7 h	4 h
9e	55 h	30 h
9f	90 h	nd
9g	37 h	nd
9h	88 h	nd
21a	64 d	nd
21b	8 d	nd
21c	52 d	nd
21d	29 d	nd
21e	100 d	nd
21f	83 h	nd
30a	17 d	nd
30b	> 70 d	nd
45	7 d	nd
46	4 d	nd
43	4 h	nd

10 nd = not determined

Abbreviations:

	Boc	t-butyloxycarbonyl
	DBU	1,3-diazabicyclo[5.4.0]undecene
	DCM	dichloromethane
5	(iv)Dde	1-(4,4-dimethyl-2,6-dioxo-cyclohexyliden)3-methyl-butyl
	DIC	diisopropylcarbodiimide
	DIEA	diisopropylethylamine
	DMAP	dimethylamino-pyridine
	DMF	N,N-dimethylformamide
10	DMSO	dimethylsulfoxide
	Dpr	diaminopropionic acid
	EDTA	ethylenediaminetetraacetic acid
	Et	ethyl
	eq	stoichiometric equivalent
15	fmoc	9-fluorenylmethoxycarbonyl
	Fmoc-Ado-OH	Fmoc-8-amino-3,6-dioxaoctanoic acid
	HFIP	hexafluoroisopropanol
	HEPES	N-(2-hydroxyethyl) piperazine-N'-(2-ethanesulfonic acid)
	HOBt	N-hydroxybenzotriazole
20	LCMS	mass spectrometry-coupled liquid chromatography
	Mal	maleimidopropionyl
	Me	methyl
	Mmt	4-methoxytrityl
	MS	mass spectrum
25	MW	molecular mass
	PyBOP	benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate
	RP-HPLC	reversed-phase high performance liquid chromatography
	RT	room temperature
30	SEC	size exclusion chromatography
	Suc	succinimidopropionyl
	TES	triethylsilane
	TFA	trifluoroacetic acid


THF	tetrahydrofurane
UV	ultraviolet
Z	benzyloxycarbonyl

CLAIMS

1. A molecule having the following structure:

EPO - Munich
60

13. Aug. 2004



5 wherein

T is D or A

D being a residue of an amine containing biologically active moiety and

A being a leaving group;

X is a spacer moiety such as R5-Y6

10 Y1, Y2 can each be either O, S, or NR6, independently of each other.

Y3, Y5 can each be either O or S, independently of each other.

Y4 is O, NR6, or -C(R7)(R8)-

Y6 is O, S, NR6, succinimide, maleimide, unsaturated carbon-carbon bonds or any heteroatom containing a free electron pair or is absent.

15 R3 is selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryl, cyano, nitro, halogen, carboxy, carboxyalkyl, alkylcarbonyl, or carboxamidoalkyl;

20 R4 is selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy, or heteroaryloxy;

25 R5 is selected from substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryl, carboxyalkyl, alkylcarbonyl or carboxamidoalkyl;

R7 and R8 are selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryls, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, cyano, or halogen;

5 R6 is selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryls;

R1 is a polymer;


10 W is selected from substituted or non-substituted linear, branched or cyclical alkyl, aryls, substituted aryls, substituted or non-substituted linear, branched or cyclical heteroalkyl, substituted or nonsubstituted heteroaryls, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy, heteroaryloxycyano, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl;

15 Nu is a nucleophile;

n is zero or a positive integer; and

Ar is a multi-substituted aromatic hydrocarbon or a multi-substituted aromatic heterocycle.

20 2. A molecule having the following structure:



wherein

T is D or A

D being a residue of an amine containing biologically active molecule and

A being a leaving group;

X is a spacer moiety such as R⁵-Y₆

Y₁, Y₂ can each be either O, S, or NR₆, independently of each other.

Y₃, Y₅ can each be either O or S, independently of each other.

5 Y₄ is O, NR₆, or -C(R₇)(R₈)-

Y₆ is O, S, NR₆, succinimide, maleimide, unsaturated carbon-carbon bonds or any heteratom containing a free electron pair or is absent

10 R₂ and R₃ are selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryl, cyano, nitro, halogen, carboxy, carboxyalkyl, alkylcarbonyl, or carboxamidoalkyl;

15 R₄ is selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy, or heteroaryloxy;

R₅ is selected from substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryl, carboxyalkyl, alkylcarbonyl or carboxamidoalkyl;

20 R₇ and R₈ are selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryl, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, cyano, or halogen;

25 R₆ is selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryl;

R₁ is a polymer;

30 W is selected from substituted or non-substituted linear, branched or cyclical alkyl, aryls, substituted aryls, substituted or non-substituted linear, branched or cyclical heteroalkyl, substituted or nonsubstituted heteroaryl, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy, heteroaryloxycyano, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl; Nu is a nucleophile;

n is zero or a positive integer; and

Ar is a multi-substituted aromatic hydrocarbon or a multi-substituted aromatic heterocycle.

- 5 3. The molecule of claim 1 or 2, wherein the biologically active moiety is selected from the group of biologically moieties consisting of small molecule biologically active agents or biopolymers.
- 10 4. The molecule of claim 3, wherein the biopolymers are selected from the group of biopolymers consisting of proteins, polypeptides, oligonucleotides and peptide nucleic acids.
- 15 5. The molecule of claim 4, wherein the polypeptides are selected from the group of polypeptides consisting of ACTH, adenosine deaminase, agalsidase, albumin, alfa-1 antitrypsin (AAT), alfa-1 proteinase inhibitor (API), alteplase, anistreplase, ancrod serine protease, antibodies (monoclonal or polyclonal, and fragments or fusions), antithrombin III, antitrypsins, aprotinin, asparaginases, biphalin, bone-morphogenic proteins, calcitonin (salmon), collagenase, DNase, endorphins, enfuvirtide, enkephalins, erythropoietins, factor VIIa, factor VIII, factor VIIla, factor IX, fibrinolysin, fusion proteins, follicle-stimulating hormones, granulocyte colony stimulating factor (G-CSF), galactosidase, glucagon, glucocerebrosidase, granulocyte macrophage colony stimulating factor (GM-CSF), phospholipase-activating protein (PLAP), gonadotropin chorionic (hCG), hemoglobins, hepatitis B vaccines, hirudin, 20 hyaluronidases, iduronidase, immune globulins, influenza vaccines, interleukins (1 alfa, 1 beta, 2, 3, 4, 6, 10, 11, 12), IL-1 receptor antagonist (rhIL-1ra), insulins, interferons (alfa 2a, alfa 2b, alfa 2c, beta 1a, beta 1b, gamma 1a, gamma 1b), keratinocyte growth factor (KGF), transforming growth factors, lactase, leuprolide, levothyroxine, luteinizing hormone, lyme vaccine, natriuretic peptide, pancrelipase, papain, parathyroid hormone, PDGF, pepsin, platelet activating factor acetylhydrolase (PAF-AH), prolactin, protein C, octreotide, secretin, sermorelin, superoxide dismutase (SOD), 25 somatropins (growth hormone), somatostatin, streptokinase, sucrase, tetanus toxin fragment, tilactase, thrombins, thymosin, thyroid stimulating hormone, 30

thyrotropin, tumor necrosis factor (TNF), TNF receptor-IgG Fc, tissue plasminogen activator (tPA), TSH, urate oxidase, urokinase, vaccines, and plant protein such as lectin and ricin.

- 5 6. The molecule of claim 4, wherein the protein is a protein prepared by recombinant DNA technology.
7. The molecule of claim 4, wherein the protein is selected from the group of proteins consisting of antibody fragments, single chain binding proteins,
10 catalytic antibodies and fusion proteins.
8. The molecule of claim 4, wherein the protein is selected from the group of proteins consisting of antibodies, calcitonin, G-CSF, GM-CSF, erythropoietins, hemoglobins, interleukins, insulins, interferons, SOD,
15 somatropin, TNF, TNF-receptor-IgC Fc.
9. The molecule of claim 3, wherein the small molecule biologically active agents are selected from the group of agents consisting of central nervous system-active agents, anti-infective, anti-neoplastic, antibacterial, anti-fungal,
20 analgesic, contraceptive, anti-inflammatory, steroidal, vasodilating, vasoconstricting, and cardiovascular agents with at least one primary or secondary amino group.
10. The molecule of claim 3, wherein the small molecule biologically active agents are selected from the group of compounds consisting of daunorubicin,
25 doxorubicin, idarubicin, mitoxantron, aminoglutethimide, amantadine, diaphenylsulfon, ethambutol, sulfadiazin, sulfamerazin, sulfamethoxazol, sulfalen, clinafloxacin, moxifloxacin, ciprofloxacin, enoxacin, norfloxacin, neomycin B, spectinomycin, kanamycin A, meropenem, dopamin, dobutamin,
30 lisinopril, serotonin, acivicin and carbutamid.
11. The molecule of claim 1 or 2, wherein R4 is selected from the group of small substituents consisting of hydrogen, methyl, ethyl, ethoxy, methoxy, and other linear alkyls, cycloalkyls or branched alkyls and heteroalkyl comprising one to


six carbon atoms.

12. The molecule of claim 1 or 2, wherein R1 is selected from the group of polymers consisting of polyalkyloxy-based polymers like poly(propylene glycol) or poly(ethylene glycol), dextran, chitosan, hyaluronic acid and derivatives, alginate, xylan, mannan, carrageenan, agarose, cellulose, starch, hydroxyethyl starch (HES) and other carbohydrate-based polymers, poly(vinyl alcohols), poly(oxazolines), poly(anhydrides), poly(ortho esters), poly(carbonates), poly(urethanes), poly(acrylic acids), poly(acrylamides) such as poly(hydroxypropylmethacrylamide) (HMPA), poly(acrylates), poly(methacrylates) like poly(hydroxyethylmethacrylate), poly(organophosphazenes), poly(siloxanes), poly(vinylpyrrolidone), poly(cyanoacrylates), poly(esters) such as poly(lactic acid) or poly(glycolic acids), poly(iminocarbonates), poly(amino acids) such as poly(glutamic acid), collagen, gelatin, copolymers, grafted copolymers, cross-linked polymers, and block copolymers from the above listed polymers.
13. The molecule of claim 1 or 2, wherein R1 is a hydrogel.
14. The molecule of claim 1 or 2, wherein R1 is a branched or hyperbranched polymer.
15. The molecule of claim 1 or 2, wherein R1 is a dendrimer or dense star polymer.
16. The molecule of claim 1 or 2, wherein R1 is a biopolymer.
17. The molecule of claim 16, wherein R1 is a protein.
18. The molecule of claim 17, wherein the protein is albumin, an antibody, fibrin, casein or any other plasma protein.


19. The molecule of any one of the above claims, wherein R1 further includes one or more biologically active substances.
- 5 20. The molecule of any one of the above claims, wherein the polymer of R1 has at least one functional group for linkage to X.
- 10 21. The molecule of claim 20, wherein the at least one functional group is selected from the group of functional groups consisting of carboxylic acid and activated derivatives, amino, maleimide, thiol, sulfonic acid and derivatives, carbonate and derivatives, carbamate and derivatives, hydroxyl, aldehyde, ketone, hydrazine, isocyanate, isothiocyanate, phosphoric acid and derivatives, phosphonic acid and derivatives, haloacetyl, alkyl halides, acryloyl, arylating agents like aryl fluorides, hydroxylamine, disulfides like pyridyl disulfide, vinyl sulfone, vinyl ketone, diazoalkanes, diazoacetyl
- 15 compounds, epoxide, oxirane, and aziridine.
22. The molecular of claim 20 or 21, wherein the at least one functional group is selected from the group of functional groups consisting of thiol, maleimide, amino, carboxylic acid and derivatives, carbonate and derivatives, carbamate and derivatives, aldehyde, and haloacetyl.
- 20
23. The molecule of one of claims 20 to 22, wherein the bond or group formed between X and R1 is selected from the group of bonds or groups consisting of disulfide, S-succinimido, amide, amino, carboxylic ester, sulphonamide, carbamate, carbonate, there, oxime, hydrazone, urea, thiourea, phosphate, phosphonate.
- 25
24. The molecule of any one of claims 20 to 23, wherein the bonds or groups formed between X and R1 is selected from the group of bonds or groups consisting of S-succinimido, amide, carbamate, and urea.
- 30
25. The molecule of any one of the above claims, wherein A is selected from the group of leaving groups consisting of chloride, bromide, fluoride, nitrophenoxy, imidazolyl, N-hydroxysuccinimidyl, N-hydroxybenzotriazolyl,

N-hydroxyazobenzotriazolyl, pentafluorophenoxy and N-hydroxysulfosuccinimidyl.

- 5 26. The molecule of claim 1 or 2, wherein the moiety $Y_4 \text{---} \overset{Y_1}{\parallel} \text{---} Y_2$ is preferably selected from the group of moieties consisting of




- 10 27. The molecule of claim 26, wherein the moiety



28. The molecule of claim 26 or 27 wherein R6 may be Nu-W.

- 20 29. The molecule of any one of claims 26 to 28 wherein Nu-W is




- 25 in which R9, R10, R11 and R12 are selected independently from hydrogen, substituted or non-substituted alkyl or heteroalkyl, substituted or non-substituted aryl or heteroaryl.

- 30 30. The molecule of claim 29, wherein R9, 10, R11 and R12 are selected independently from hydrogen or substituted or non-substituted alkyl.

31. The molecule of any of the above claims, wherein Nu is selected from the group of nucleophiles consisting of primary, secondary and tertiary amino groups, thiol, carboxylic acid, hydroxylamine and hydrazine.

5

10



20

25

30


36. Use of the molecule of any one of the above claims 1 to 35, in which T is a leaving group A, for covalent conjugation with a biologically active moiety and linkage with a carrier.

37. The use of claim 36, wherein the biologically active moiety is an amine-containing molecule.

38. The use of claim 36 or 37, wherein the biologically active moiety is a peptide, polypeptide or protein.

39. Method for the synthesis of a polymeric prodrug comprising:

- providing a starting molecule of Formula II



- synthesising at least one intermediate compound from the starting molecule of Formula II; and

- attaching an amine-containing biologically active moiety D to the at least one intermediate compound to form the polymeric prodrug;

wherein

Y₂ is selected from O, S, or NR₆

Y₃ is selected from O or S

X is a spacer moiety such as R₅-Y₆;

R₃ is selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl, cyano, nitro, halogen, carboxy, carboxyalkyl, alkylcarbonyl or carboxamidoalkyl;

R₄ is selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy or heteroaryloxy;

R₅ is selected from substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl;


R₆ is selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl;

Y_6 is O, S, NR6, succinimide, maleimide, unsaturated carbon-carbon bonds or a heteratom containing a free electron pair;


n is zero or a positive integer and

Ar is a multi-substituted aromatic hydrocarbon or a multi-substituted aromatic heterocycle.

40. The method of claim 39, wherein a first one of the at least one intermediate compounds is an intermediate molecule of Formula III



synthesised by acylating Y_2 with



and attaching a first protecting group PG_1 wherein Nu is a nucleotide;

W is selected from substituted or non-substituted linear, branched or cyclical alkyl, aryls, substituted aryls, substituted or non-substituted linear, branched or cyclical heteroalkyl, substituted or nonsubstituted heteroaryl, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkoxy, aryloxy,

heteroaryloxy, cyano, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl;


Y_1 is selected from the group of moieties consisting of O, S, or NR6;

Y_4 is selected from the group of moieties consisting of, or $-C(R7)(R8)-O$, NR6;

R7 and R8 are selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, cyano or halogen.

PG_1 is a first protecting group or absent.

41. The method of claim 40, wherein a second one of the at least one intermediate compound is an intermediate compound of formula IV



formed by activating the compound of Formula III with an activating agent,
wherein A is a leaving group and Y₅ is selected from O or S.

42. The method of claim 41, wherein A is selected from chloride, bromide, fluoride, nitrophenoxy, imidazolyl, N-hydroxysuccinimidyl, N-hydroxybenzotriazolyl, N-hydroxyazobenzotriazolyl, pentafluorophenoxy and N-hydroxysulfosuccinimidyl.

43. The method of one of claims 41 or 42, wherein the activating agent is selected from the group of activating agents consisting of 4-nitrophenyl chloroformate or disuccinyl carbonate.


44. The method of one of claims 41 or 42, wherein the amine-containing biologically active moiety D is attached to the second one of the at least one intermediary compound (Formula IV) by displacement of the leaving group A.

45. The method of claim 44 comprising a step of removal of the reversible first protecting group PG₁ from the second one of the at least one intermediate compound.


46. The method of claim 45, wherein the step of removal of the reversible protection group PG₁ is carried out using a reagent selected from the group of reagents consisting of trifluoroacetic acid or DTT.

47. The method of any one of claims 45 or 46, further comprising a step of attaching a polymer R1 to X.

48. The method of claim 40, further comprising a step of removing the first protection group PG₁ and attaching a polymer R1 to X to form a third one of the at least one intermediate compound and having a Formula VI.



49. The method of claim 48, further comprising a step of activating the third one
of the at least one intermediate compound by using an activating agent to form
a fourth one of the at least one intermediate compound and having a Formula
VII and wherein A is a leaving group



50. The method of claim 49, wherein A is selected from chloride, bromide,
fluoride, nitrophenoxy, imidazolyl, N-hydroxysuccinimidyl, N-
hydroxybenzotriazolyl, N-hydroxyazobenzotriazolyl, pentafluorophenoxy and
N-hydroxysulfosuccinimidyl.

51. The method of claim 49 or 50, wherein the activating agent is selected from
the group of activating agents consisting of 4-nitrophenyl chloroformate or
disuccinyl carbonate.


52. The method of one of claim 49 to 51, wherein the amine-containing
biologically active moiety D is attached to the compound of Formula VII by
displacement of the leaving group A.

53. The method of claim 39, further comprising a step of reacting the starting
molecule of Formula II with a first polymer R1 to attach the first polymer R1
to X.

54. The method of claim 53, further comprising a step of protecting Y₂ with a
second protective group PG₂.

55. The method of claim 53 or 54, further comprising the step of activating the
molecule with an activating agent and reacting with an amine-containing
biologically active moiety D to form a fifth one of the at least one intermediate

5



10

$$\text{Nu}-\text{W}-\text{Y}_4-\overset{\text{Y}_1}{\parallel}$$

15

20

25

30

Page 14/21

5



15

20

25

30


$$\text{Nu}-\text{W}-\text{Y}_4 \begin{array}{l} \text{Y} \\ \parallel \\ \text{Y}_1 \end{array}$$

Page 15/21

Y1 is selected from the group of moieties consisting of O, S, or NR6;
 Y4 is selected from the group of moieties consisting of -C(R7)(R8)-; O, or
 NR6; and

5 R7 and R8 are selected from hydrogen, substituted or non-substituted linear,
 branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or
 non-substituted heteroaryl, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl,
 cyano or halogen.

10 65. The method of claim 62, wherein the second protecting group PG₂ is removed
 from Y₂ and Y₂ is acylated with




15 wherein W is selected from substituted or non-substituted linear, branched or
 cyclical alkyl, aryls, substituted aryls, substituted or non-substituted linear,
 branched or cyclical heteroalkyl, substituted or nonsubstituted heteroaryls,
 substituted or non-substituted linear, branched, or cyclical alkoxy, substituted
 or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy,
 heteroaryloxycyano, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl;
 Nu is a nucleotide;

20 Y1 is selected from the group of moieties consisting of O, S, or NR6;
 Y4 is selected from the group of moieties consisting of -C(R7)(R8)- O, or
 NR6; and
 R7 and R8 are selected from hydrogen, substituted or non-substituted linear
 alkyl, branched alkyl or cyclical alkyl, aryl, substituted aryl, substituted or
 non-substituted heteroalkyl or heteroaryl, carboxyalkyl, alkylcarbonyl,
 25 carboxamidoalkyl, cyano or halogen.


66. The method of claim 65 comprising a step of removing the first protecting
 group PG₁ from X and attaching a polymer R1.

30 67. The method of claim 43 comprising a step acylating Y₂ with



- to form a third one of the at least one intermediate compound;
wherein W is selected from substituted or non-substituted linear, branched or
cyclical alkyl, aryls, substituted aryls, substituted or non-substituted linear,
branched or cyclical heteroalkyl, substituted or nonsubstituted heteroaryls,
5 substituted or non-substituted linear, branched, or cyclical alkoxy, substituted
or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy,
heteroaryloxycyano, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl;
Nu is a nucleotide;
Y₁ is selected from the group of moieties consisting of O, S, or NR₆;
10 Y₄ is selected from the group of moieties consisting of -C(R₇) (R₈)-O or
NR₆; and
R₇ and R₈ are selected from hydrogen, substituted or non-substituted linear
alkyl, branched alkyl or cyclical alkyl, aryl, substituted aryl, substituted or
non-substituted heteroalkyl or heteroaryl, carboxyalkyl, alkylcarbonyl,
15 carboxamidoalkyl, cyano or halogen.


68. The method of claim 67, further comprising a step of activating the third one
of the at least one intermediate compound by using an activating agent to form
a seventh one of the at least one intermediate compound and having a Formula
20 VII and wherein A is a leaving group



- 25 69. The method of claim 68, wherein A is selected from the group of leaving
groups consisting of chloride, nitrophenoxy, imidazolyl, N-
hydroxysuccinimidyl, N-hydroxybenzotriazolyl, N-hydroxyazobenzotriazolyl,
pentafluorophenoxy and N-hydroxysulfosuccinimidyl.
- 30 70. The method of claim 68 or 69, wherein the activating agent is selected from
the group of activating agents consisting of 4-nitrophenyl chloroformate or
disuccinyl carbonate.

71. The method of one of claims 68 to 70, wherein the amine-containing biologically active moiety D is attached to the compound of Formula VII by displacement of the leaving group A.

5 72. The method of any one of claims 39 to 71 wherein the starting molecule of Formula II is replaced by a starting molecule of Formula IIb



10

wherein R2 is selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyls or heteroalkyls, aryls, substituted or non-substituted heteroaryl, cyano, nitro, halogen, carboxy, carboxyalkyl, alkylcarbonyl or carboxamidoalkyl.

15

73. A method for hydrolysing the molecule of any one of claims 1 to 35 comprising a step of placing the molecule in solution with a pH of approximately 7.4

20

74. The method of claim 73, wherein the solution is an extra-cellular fluid.

75. Method of administration of an amine-containing moiety to a living organism comprising:

25

- a first step of attaching the amine-containing moiety through an amino group to a carrier molecule by means of a nucleophile-containing linker;
- a second step of administering the amine-containing moiety and the carrier molecule to the living organism; and
- a third step of cleaving the amine-containing moiety from the carrier molecule by means of a substantially non-enzymatic reaction.

30

76. The method of claim 75, wherein the amine-containing moiety is a biologically active moiety.

77. The method of claim 76 wherein the biologically active moiety is selected from the group of biologically moieties consisting of small molecule biologically active agents or biopolymers.
- 5 78. The molecule of claim 76, wherein the biopolymers are selected from the group of biopolymers consisting of proteins, polypeptides, oligonucleotides and peptide nucleic acids.
- 10 79. The method of claim 78, wherein the polypeptides are selected from the group of polypeptides consisting of ACTH, adenosine deaminase, agalsidase, albumin, alfa-1 antitrypsin (AAT), alfa-1 proteinase inhibitor (API), alteplase, anistreplase, anrod serine protease, antibodies (monoclonal or polyclonal, and fragments or fusions), antithrombin III, antitrypsins, aprotinin, asparaginases, biphalin, bone-morphogenic proteins, calcitonin (salmon), collagenase, DNase, endorphins, enfuvirtide, enkephalins, erythropoietins, factor VIIa, factor VIII, factor VIIa, factor IX, fibrinolysin, fusion proteins, follicle-stimulating hormones, granulocyte colony stimulating factor (G-CSF), galactosidase, glucagon, glucocerebrosidase, granulocyte macrophage colony stimulating factor (GM-CSF), phospholipase-activating protein (PLAP), gonadotropin chorionic (hCG), hemoglobins, hepatitis B vaccines, hirudin, hyaluronidases, iduronidase, immune globulins, influenza vaccines, interleukins (1 alfa, 1 beta, 2, 3, 4, 6, 10, 11, 12), IL-1 receptor antagonist (rhIL-1ra), insulins, interferons (alfa 2a, alfa 2b, alfa 2c, beta 1a, beta 1b, gamma 1a, gamma 1b), keratinocyte growth factor (KGF), transforming growth factors, lactase, leuprolide, levothyroxine, luteinizing hormone, lyme vaccine, natriuretic peptide, pancrelipase, papain, parathyroid hormone, PDGF, pepsin, platelet activating factor acetylhydrolase (PAF-AH), prolactin, protein C, octreotide, secretin, sermorelin, superoxide dismutase (SOD), somatropins (growth hormone), somatostatin, streptokinase, sucrase, tetanus toxin fragment, tilactase, thrombins, thymosin, thyroid stimulating hormone, thyrotropin, tumor necrosis factor (TNF), TNF receptor-IgG Fc, tissue plasminogen activator (tPA), TSH, urate oxidase, urokinase, vaccines, and plant protein such as lectin and ricin.
- 15 20 25 30

80. The method of claim 78, wherein the protein is a protein prepared by recombinant DNA technology.

5 81. The method of claim 78, wherein the protein is selected from the group of proteins consisting of antibody fragments, single chain binding proteins, catalytic antibodies and fusion proteins.

10 82. The method of claim 77, wherein the small molecule biologically active agents are selected from the group of agents consisting of central nervous system-active agents, anti-infective, anti-neoplastic, antibacterial, anti-fungal, analgesic, contraceptive, anti-inflammatory, steroidal, vasodilating, vasoconstricting, and cardiovascular agents with at least one primary or secondary amino group.

15 83. The method of claim 77, wherein the small molecule biologically active agents are selected from the group of compounds consisting of daunorubicin, doxorubicin, idarubicin, mitoxantron, aminoglutethimide, amantadine, diaphenylsulfon, ethambutol, sulfadiazin, sulfamerazin, sulfamethoxazol, sulfalen, clinafloxacin, moxifloxacin, ciprofloxacin, enoxacin, norfloxacin,
20 neomycin B, spectinomycin, kanamycin A, meropenem, dopamin, dobutamin, lisinopril, serotonin, acivicin, and carbutamid.

25 84. The method of one of claims 75 to 83, wherein the third step is carried out in an extra-cellular fluid.

85. The method of one of claims 75 to 84, wherein the substantially non-enzymatic reaction comprises a step of hydrolysis.


30 86. The method of one of claims 75 to 85, wherein the substantially non-enzymatic reaction comprises a step of intramolecular cyclization.

87. The method of one of claims 75 to 86, further comprising a step of sterically protecting at least part of the linker by a sterically demanding carrier.


- 5
88. In an amine-containing moiety attached to a carrier by a nucleophile-containing linker, a method of cleaving the amine-containing moiety from the carrier by a substantially non-enzymatic reaction of the nucleophile-containing linker.
89. The method of claim 88, wherein the substantially non-enzymatic reaction is carried out at a pH of approximately 7.4.
- 10
90. The method of claim 88 or 89, wherein the amine-containing moiety attached to the carrier is cleaved in an extra-cellular fluid.
91. The method of one of claims 88 to 90, wherein the substantially non-enzymatic reaction comprises a step of hydrolysis.
- 15
92. The method of one of claims 88 to 91, wherein the substantially non-enzymatic reaction comprises a step of intramolecular cyclization.
- 20
93. The method of one of claims 88 to 92, further comprising a step of sterically protecting at least part of the nucleophile-containing linker by a sterically demanding carrier.
94. The method of any one of claims 88 to 93, wherein the amine-containing moiety is a biologically active moiety.

ABSTRACT 13. Aug. 2004

The invention relates to a molecule having the following structure:



5 or



for use as a polymeric prodrug.

EPO - Munich
60
13. Aug. 2004

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8





Fig. 9






Fig. 11

